Acknowledgement

The present thesis is the collection of the studies which have been carried out under the direction of Prof. Dr. Hiroyuki Nishide, Department of Applied Chemistry in Waseda University, during the 2000-2002. The author expresses the greatest acknowledgement to Prof. Dr. Hiroyuki Nishide for his invaluable suggestion, discussion, and continuous encouragement throughout this work.

The author expresses her sincere gratitude to Honorary Prof. Dr. Eishun Tsuchida and Assoc. Prof. Shinji Takeoka for their invaluable advices comments as well as continuous encouragement for this study.

The author wishes to thank Prof. Dr. Kiyotaka Sakai for his efforts as members on judging committee for the doctoral thesis.

The author express the special thank to all collaborators, Assoc. Prof. Dr. Baoqing Shentu (Zhejing Univ.), Mr. Takayuki Arai, Mr. Hirotsugu Araihara, and Mr. Hirokazu Shibata, for their assistance in the experimental work. The author deeply thanks all the members in the laboratory for their kind assistance.

The author would like to thank Dr. Yong Soo Kang (KIST), Prof. Dr. Jongok Won (Sejong Univ.), and Dr. Jong Hak Kim (Yonsei Univ.) for their productive comments and unlimited helpful discussion on this study.

The author would like to thank Prof. Dr. Ichiro Okura (Tokyo Institute of Technology), Assoc. Prof. Dr. Yutaka Amao (Oita Univ.), and Dr. Aiko Nakao (RIKEN) for many helpful suggestions and productive comments during the course of this work.

Finally, the author express the deepest gratitude to her grandparents, especially Honorary Prof. Isao Shinohara (Waseda Univ.), my parents Mr. Satoshi Shinohara, Mrs. Yasuko Shinohara, and my brothers, Mr. Susumu Shinohara and Mr. Takashi Shinohara, and Yoshiyuki Tanaka, for their endless love and kindness support.

December, 2002 Hiromi Shinohara