早稲田大学審査学位論文(博士)

POLYNOMIAL INVARIANTS OF PERIODIC KNOTS

(周期的な結び目の多項式不変量)

BY YOSHIYUKI YOKOTA

A dissertation submitted for the degree of
Doctor of Science at Waseda University

October 1992

POLYNOMIAL INVARIANTS OF PERIODIC KNOTS

BY YOSHIYUKI YOKOTA

A dissertation submitted for the degree of
Doctor of Science at Waseda University
October 1992

ACKNOWLEDGEMENTS

I am so grateful to Professor Shin'ichi Suzuki for his valuable suggestions, comments and encouragements at Waseda university.

YOSHIYUKI YOKOTA
DEPARTMENT OF MATHEMATICS
KYUSHU UNIVERSITY
FUKUOKA, 812
JAPAN

CONTENTS

INTRODUCTION

CHAPTER 1. JONES POLYNOMIALS OF PERIODIC KNOTS

- 1.1. MAIN THEOREMS.
- 1.2. TEMPERLEY-LIEB ALGEBRAS.
- 1.3. PROOF OF THEOREMS.
- 1.4. EXAMPLES AND REMARKS.

CHAPTER 2. SKEIN POLYNOMIALS OF PERIODIC KNOTS

- 2.1. MAIN THEOREM.
- 2.2. HECKE ALGEBRAS.
- 2.3. TORUS LINKS.
- 2.4. TORUS KNOTS.
- 2.5. PERIODIC REDUCTIONS.

CHAPTER 3. KAUFFMAN POLYNOMIALS OF PERIODIC KNOTS

- 3.1. MAIN THEOREM.
- 3.2. BRAUER'S ALGEBRAS.
- 3.3. BIRMAN-WENZL-MURAKAMI ALGEBRAS.
- 3.4. TORUS LINKS.
- 3.5. TORUS KNOTS.
- 3.6. PERIODIC REDUCTIONS.

CHAPTER 4. A RELATIONSHIP BETWEEN SKEIN AND KAUFFMAN POLYNOMIALS

4.1. MAIN THEOREM.

4.2. TORUS KNOTS.

4.3. PERIODIC REDUCTIONS.

REFERENCES

A LIST OF PAPERS BY YOSHIYUKI YOKOTA

POLYNOMIAL INVARIANTS OF PERIODIC KNOTS

INTRODUCTION

We consider “knots”, which are smoothly embedded circles in 3-sphere S^3 . The theory of knots is a beautiful subject in the low dimensional topology. In fact, knots often show us various interesting phenomena, and provide us a lot of mathematical topics.

Among such ones, in this paper, our interest will be restricted to beautiful symmetries of knots. Some knots show geometric symmetries, which are called “symmetric” or “periodic” knots. A topological description of symmetries of knots can be given as follows. A knot K in S^3 will be said to have period r , if K is mapped onto itself by an orientation preserving homeomorphism f of S^3 which has period r with the axis $\text{Fix } f$ away from K . By the positive solution of “Smith conjecture”[25], f is a standard rotation of S^3 with unknotted axis $\text{Fix } f$. Thus, the quotient space of f also becomes 3-sphere Σ^3 and so, the image of K will be a new knot in Σ^3 . This will be called a “factor knot” of K .

A KNOT WITH PERIOD 3

Various tools in knot theory have been applied to this topic. It seems that there has been a common policy:

“Symmetries of knots should be reflected in their algebraic invariants.”

In this direction, representations of knot groups[23,38], homology of covering spaces[10,29], signatures[9], Alexander polynomials[6,29] and hyperbolic structures[1] of knots has been often explored, and these work enough to determine symmetries of knots with at most 10 crossings, except several knots[18]. For instance, due to [29], we now describe a useful result on Alexander polynomials of periodic knots.

Let r be a prime. Note that a knot has prime period whenever she has certain period, and so we will keep this assumption throughout of this paper. Let K be a knot with period r and k its factor knot. A relationship of Alexander polynomials $\Delta_K(t)$ and $\Delta_k(t)$ is given by

$$\Delta_K(t) \equiv (1 + t + \cdots + t^{r-1})^{r-1} \Delta_k(t)^r \pmod{r},$$

where λ denotes the linking number of K and the axis of rotation.

In 1984, V.F.R. Jones introduced a new polynomial invariant in knot theory[12]. This new polynomial, which is now called "Jones polynomials", is not only powerful to distinguish knots, but also useful enough to settle classical conjectures in knot theory[31,32].

After the appearance of Jones polynomials, several authors quickly discovered a new polynomial invariant[8], that is "Skein polynomials", which is a 2 variable generalization of Jones. Furthermore, L.H. Kauffman introduced a concept "regular isotopy" of knots and defined another polynomial invariant, so called "Kauffman polynomials", using his regular isotopy invariant of knots[17].

K. Murasugi[33] first applied Jones polynomials to the study of periodic knots. He described a relationship between Jones polynomials of a periodic knot and its factor knot, which is analogous to his own work on Alexander polynomials, and showed that 10_{105} in [35], one of the undecided cases, has no period. After this work, P. Traczyk[39] found a certain condition of Jones polynomials for periodic knots and also proved that 10_{101} in [35] can not have any period. Of course, other polynomials have been also applied to the study of periodic knots. J.H. Przytycki[34] started the study of Skein and Kauffman polynomials for periodic knots and have given several criteria which are analogous to Jones'. Furthermore, P. Traczyk[40,41] found an interesting regularity of Skein polynomials, which works well even for knots with small period, and could prove several undecided knots have no period.

Thus, new polynomials seems effective on the study of periodic knots. However, present results from this point of view does not look like satisfactory. They indeed look like having further informations on symmetries of knots. So, in this paper, we will investigate polynomial invariants for periodic knots in our own way, and will discover mysterious behaviours of them. Indeed, symmetries of knots are amazingly reflected in their polynomials, and perhaps, no one could expect such phenomena.

We begin with Jones polynomials in Chapter 1. Results obtained there will be improvements of K. Murasugi[33] and P. Traczyk[39]. In Chapter 2, we will introduce Skein polynomials and completely settle a conjecture by P. Traczyk. Unexpectedly strong regularity of Skein polynomials will be shown there. In Chapter 3, we will introduce Kauffman polynomials and establish a regularity similar to Chapter 2. Kauffman polynomials of torus knots will be explicitly written down there, and play an important role in the proof of Theorem.

Results which we will obtain in Chapters 2 and 3 will suggest a curious correspondence of Skein and Kauffman polynomials for periodic knot, while it is known these 2 invariants are entirely independent in general. Chapter 4 will be devoted to explain this curious correspondence and to formulate in a suitable way, that is, by defining a new polynomial invariant.

Unfortunately, we had to exclude knots with period 2 from our concern. In fact, polynomial invariants of knots with period 2 look like having no specific feature. It

might be hard to exhibit an effective regularity for knots with period 2 using polynomial invariants. No one has succeeded this at present stage. So, throughout of this paper, we further assume r denotes an odd prime.

Examples of polynomials reflecting geometric information of knots, as symmetries, are scarcely known and so, results exhibited in this paper will be meaningful as such one.

I now think that 2 dimensional informations of knots, for example, minimal number of crossings, braid index, alternativity, positivity,... tends to be reflected in their polynomials while 3 or 4 dimensional informations of knots, for example, unknotting number, bridge index, hyperbolicity, sliceness,... tends to be carelessly ignored. On symmetries of knots, the positive solution of Smith conjecture fortunately enables us to treat periodic knots in rather 2 dimensional way, and polynomial invariants have indeed caught them.

I now expect these mysterious effects of symmetries of knots appearing in their polynomial invariants would give a clue for establishing a geometric characterization of new polynomial invariants, which would be most mysterious.

CHAPTER 1. JONES POLYNOMIALS OF PERIODIC KNOTS

In this Chapter, we will study Jones polynomials of knots with period r . The positive solution of Smith conjecture enables us to treat periodic knots combinatorially. K. Murasugi[33] first applied Jones polynomials to the study of periodic knots, which uses clever combinatorics from Jones polynomials.

We shall use an algebra coming from Jones polynomials and describe symmetries of knots in terms of this algebra.

1.1. MAIN THEOREMS.

Let us first define "Jones polynomials".

Definition[12]. The *Jones polynomial* $V_L(t) \in Z[t^{\pm \frac{1}{2}}]$ of an oriented link L is uniquely defined by

$$V_{\bigcirc}(t) = 1, \\ t^{-1} V_{\text{cross}}(t) - t V_{\text{uncross}}(t) = (t^{\frac{1}{2}} - t^{-\frac{1}{2}}) V_{\text{parallel}}(t).$$

We will show the followings in this Chapter.

THEOREM 1.1. *Let K be a knot with period r .*

(i) *If $\text{lk}(K, \text{Fix } f)$ is odd, then*

$$V_K(t) - V_K(t^{-1}) \equiv 0 \pmod{(r, t^{2r} - 1)}.$$

(ii) *If $\text{lk}(K, \text{Fix } f)$ is even, then*

$$V_K(t) - V_K(t^{-1}) \equiv 0 \pmod{(r, t^r - 1)},$$

$$V_K(t) + V_K(t^{-1}) \equiv 0 \pmod{(r, \frac{t^r + 1}{t + 1})}.$$

Here, f is a rotation of S^3 preserving K .

THEOREM 1.2. *Let K be a knot with period r . We shall denote the factor knot of K by k .*

(i) *If $\text{lk}(K, \text{Fix } f)$ is odd, then*

$$V_K(t) \equiv V_k(t)^r \pmod{(r, (t^{r+1} - 1)(t^{r-1} - 1))}.$$

(ii) *If $\text{lk}(K, \text{Fix } f)$ is even, then*

$$V_K(t) \equiv (t^{\frac{1}{2}} + t^{-\frac{1}{2}})^{r-1} V_k(t)^r \pmod{(r, \frac{(t^{r+1} - 1)(t^{r-1} - 1)}{t + 1})}.$$

In the case of knots, the following is a direct consequence of Theorem 1.1.

THEOREM 1.3[39]. If K is a knot with period r ,

$$V_K(t) - V_K(t^{-1}) \equiv 0 \pmod{(r, t^r - 1)}.$$

Theorem 1.2 also generalizes Murasugi[33].

THEOREM 1.4[33]. Under the same assumption as in Theorem 1.2,

$$V_K(t) \equiv V_k(t)^r \pmod{\begin{cases} (r, (t+1)^{2(r-1)} - t^{r-1}) & \text{if } \text{lk}(K, \text{Fix } f) \text{ is odd,} \\ (r, (t+1)^{r-1} - t^{\frac{r-1}{2}}) & \text{if } \text{lk}(K, \text{Fix } f) \text{ is even.} \end{cases}}$$

The proof of Theorem 1.1 and Theorem 1.2 will be given in section 1.3. We shall prepare several materials from a view point of representation theory in the following section.

1.2. TEMPERLEY-LIEB ALGEBRAS.

We will first introduce "Temperley-Lieb algebras". Jones polynomials will be then described as a weighted sum of irreducible characters of this algebras.

Definition. Fix a complex number $A \neq 0$ and a positive integer n . Then, *Temperley-Lieb algebra* J_n is defined as an algebra over $\mathbb{C}$ with generators

$$1, e_1, e_2, \dots, e_{n-1}$$

and relations

$$e_i^2 = -(A^2 + A^{-2})e_i,$$

$$e_i e_{i \pm 1} e_i = e_i,$$

$$e_i e_j = e_j e_i \quad \text{if } |i - j| > 1.$$

As is well-known, J_n is semisimple when A is not a root of unity and decomposes into $[n/2] + 1$ simple components. Let $\rho_{n,i}$'s denote irreducible representations of J_n and $\chi_{n,i}$'s their characters, where $0 \leq i \leq [n/2]$.

To connect periodic knots and Temperley-Lieb algebras, we shall introduce

Definition[26]. Let F_n be a semigroup generated by tangles with n strings given by

ϵ_i
 σ_i
 σ_i^{-1}

where the product $\xi\eta$ of $\xi, \eta \in F_n$ will be defined by joining n points at the bottom of ξ with n points at the top of η . As braids, we can define the closure of ξ , denoted by $\hat{\xi}$, naturally.

Let $\pi_n : F_n \rightarrow J_n$ a semigroup homomorphism defined by

$$\begin{aligned}\pi_n(\epsilon_i) &= e_i, \\ \pi_n(\sigma_i) &= A^{-1} + Ae_i, \\ \pi_n(\sigma_i^{-1}) &= A + A^{-1}e_i.\end{aligned}$$

In the proof of Theorems, we will essentially use the following, which gives a character formula for Kauffman's bracket [16].

THEOREM 1.5 [27]. For $\xi \in F_n$, we have

$$\langle \hat{\xi} \rangle = \sum_{i=0}^{[n/2]} a_{n,i}(A) \chi_{n,i}(\pi_n(\xi)),$$

where $\langle \hat{\xi} \rangle$ denotes Kauffman's bracket of $\hat{\xi}$ and

$$a_{n,i}(A) = (-1)^{n+1} \frac{A^{2n+2-4i} - A^{-2n-2+4i}}{A^4 - A^{-4}}.$$

1.3 PROOF OF THEOREMS.

Let K be a knot with period r and k its factor knot. Then, the following proposition may be obvious.

PROPOSITION 1.6. There exists n and $\xi \in F_n$ such that K and k are closures of ξ^r and ξ respectively. Furthermore, n is odd or even according as $\text{lk}(K, \text{Fix } f)$ is odd or even.

Orientations on K and k naturally induce orientations on $\hat{\xi}^r$ and $\hat{\xi}$. So, by $w(K)$ and $w(k)$, we shall denote algebraic crossing numbers of $\hat{\xi}^r$ and $\hat{\xi}$ respectively.

+1

-1

Note that $w(K) = rw(k)$ and Jones polynomials of $\hat{\xi}^r$ and $\hat{\xi}$ are given by

$$\begin{aligned}V_K(A^4) &= (-A^3)^{w(K)} \langle \hat{\xi}^r \rangle, \\ V_k(A^4) &= (-A^3)^{w(k)} \langle \hat{\xi} \rangle.\end{aligned}$$

We now expand right hand sides by using Theorem 1.5. Then,

$$\begin{aligned}
V_K(A^4) &= (-A^3)^{w(K)} \langle \hat{\xi}^r \rangle \\
&= \sum_{i=0}^{[n/2]} a_{n,i}(A) (-A^3)^{rw(k)} \chi_{n,i}(\pi_n(\xi^r)), \\
V_k(A^4) &= (-A^3)^{w(k)} \langle \hat{\xi} \rangle \\
&= \sum_{i=0}^{[n/2]} a_{n,i}(A) (-A^3)^{w(k)} \chi_{n,i}(\pi_n(\xi)).
\end{aligned}$$

Here, we put $f_i(A) = (-A^3)^{w(k)} \chi_{n,i}(\pi_n(\xi)) \in Z[A^{\pm 1}]$.

LEMMA 1.7.

$$(-A^3)^{rw(k)} \chi_{n,i}(\pi_n(\xi^r)) \equiv f_i(A^r) \pmod{r}.$$

Proof. As $\rho_{n,i}(\pi_n(\xi))$'s are matrices over $Z[A^{\pm 1}]$, we have

$$\begin{aligned}
\chi_{n,i}(\pi_n(\xi^r)) &= \text{tr}(\rho_{n,i}(\pi_n(\xi^r))) \\
&= \text{tr}(\rho_{n,i}(\pi_n(\xi))^r) \\
&\equiv \text{tr}(\rho_{n,i}(\pi_n(\xi)))^r \pmod{r}.
\end{aligned}$$

That is,

$$\chi_{n,i}(\pi_n(\xi^r)) \equiv \chi_{n,i}(\pi_n(\xi))^r \pmod{r}.$$

Thus, we obtain

$$\begin{aligned}
(-A^3)^{rw(k)} \chi_{n,i}(\pi_n(\xi^r)) &\equiv (-A^3)^{w(k)} \chi_{n,i}(\pi_n(\xi))^r \\
&\equiv f_i(A)^r \\
&\equiv f_i(A^r) \pmod{r}. \quad \square
\end{aligned}$$

We have to consider 2 cases of Theorems.

Case (i). $\text{lk}(K, \text{Fix } f)$ is odd.

By Proposition 1.6, n is odd and $a_{n,i}(A)$'s are in $Z[A^{\pm 4}]$ from Theorem 1.5. Decompose $f_i(A)$'s into

$$f_i(A) = g_i(A) + h_i(A),$$

where $g_i(A)$'s are in $Z[A^{\pm 4}]$ and $h_i(A)$'s are in $Z[A^{\pm 1}] - Z[A^{\pm 4}]$. Clearly, such a decomposition is unique. By Lemma 1.7, we have

$$V_K(A^4) \equiv \sum_{i=0}^{[n/2]} a_{n,i}(A) g_i(A^r) + \sum_{i=0}^{[n/2]} a_{n,i}(A) h_i(A^r) \pmod{r}$$

and

$$V_k(A^\pm) = \sum_{i=0}^{[n/2]} a_{n,i}(A)g_i(A) + \sum_{i=0}^{[n/2]} a_{n,i}(A)h_i(A).$$

As K and k are knots, not links, $V_K(A^\pm)$ and $V_k(A^\pm)$ are in $Z[A^{\pm 4}][12]$. This implies

$$\sum_{i=0}^{[n/2]} a_{n,i}(A)h_i(A^r) \equiv \sum_{i=0}^{[n/2]} a_{n,i}(A)h_i(A) \equiv 0 \pmod{r}.$$

Since $a_{n,i}(A) = a_{n,i}(A^{-1})$, we have

$$V_K(A^\pm) - V_K(A^{\mp}) \equiv \sum_{i=0}^{[n/2]} a_{n,i}(A)(g_i(A^r) - g_i(A^{-r})) \pmod{r}$$

and $A^{4r} - A^{-4r}$ divides the right hand side because $g_i(A^r)$'s are in $Z[A^{\pm 4r}]$. By replacing t for $A^\pm$, we obtain Theorem 1.1(i).

Furthermore, we can write

$$V_K(A^\pm) - V_k(A^\pm)^r \equiv \sum_{i=0}^{[n/2]} (a_{n,i}(A) - a_{n,i}(A)^r)g_i(A^r) \pmod{r}.$$

Since

$$a_{n,i}(A) = A^{2n-2-4i} + A^{2n-10-4i} + \dots + A^{-2n+10+4i} + A^{-2n+2+4i},$$

it is easy to show

$$a_{n,i}(A) - a_{n,i}(A)^r \equiv 0 \pmod{(r, (A^{4(r+1)} - 1)(A^{4(r-1)} - 1))}.$$

In fact,

$$\begin{aligned} A^{4m} + A^{-4m} - (A^{4m} + A^{-4m})^r &\equiv -A^{-4mr}(A^{4m(r+1)} - 1)(A^{4m(r-1)} - 1) \\ &\equiv 0 \pmod{(r, (A^{4(r+1)} - 1)(A^{4(r-1)} - 1))}. \end{aligned}$$

This gives Theorem 1.2(i).

Case (ii). $\text{lk}(K, \text{Fix } f)$ is even.

By Proposition 1.6, n is even. Theorem 1.5 tells us that $(A^2 + A^{-2})a_{n,i}(A)$'s are in $Z[A^{\pm 4}]$ while $a_{n,i}(A)$'s are never in $Z[A^{\pm 4}]$. Then, by an argument similar to that of Case (ii), we obtain

$$\begin{aligned} (A^2 + A^{-2})V_K(A^\pm) &\equiv \sum_{i=0}^{[n/2]} (A^2 + A^{-2})a_{n,i}(A)y_i(A^r) \pmod{r}, \\ (A^2 + A^{-2})V_k(A^\pm) &= \sum_{i=0}^{[n/2]} (A^2 + A^{-2})a_{n,i}(A)y_i(A), \end{aligned}$$

where $y_i(A)$'s are in $Z[A^{\pm 2}] - Z[A^{\pm 4}]$ and so,

$$\begin{aligned}
& (A^2 + A^{-2})(V_K(A^4) - V_K(A^{-4})) \\
& \equiv \sum_{i=0}^{[n/2]} (A^2 + A^{-2})a_{n,i}(A)(y_i(A^r) - y_i(A^{-r})) \pmod{r}, \\
& (A^2 + A^{-2})(V_K(A^4) + V_K(A^{-4})) \\
& \equiv \sum_{i=0}^{[n/2]} (A^2 + A^{-2})a_{n,i}(A)(y_i(A^r) + y_i(A^{-r})) \pmod{r}.
\end{aligned}$$

Since $y_i(A^r)$'s are in $Z[A^{\pm 2r}] - Z[A^{\pm 4r}]$, $y_i(A^r) - y_i(A^{-r})$'s and $y_i(A^r) + y_i(A^{-r})$'s are divisible by $A^{2r} - A^{-2r}$ and $A^{2r} + A^{-2r}$ respectively. By replacing t for A^4 , we obtain Theorem 1.1(ii).

Furthermore,

$$\begin{aligned}
& (A^2 + A^{-2})V_K(A^4) - [(A^2 + A^{-2})V_K(A^4)]^r \\
& \equiv \sum_{i=0}^{[n/2]} ((A^2 + A^{-2})a_{n,i}(A) - [(A^2 + A^{-2})a_{n,i}(A)]^r)y_i(A^r) \pmod{r}.
\end{aligned}$$

As in Case (i), it is also proved

$$(A^2 + A^{-2})a_{n,i}(A) - [(A^2 + A^{-2})a_{n,i}(A)]^r \equiv 0 \pmod{(r, (A^{4r+4} - 1)(A^{4r-4} - 1))}.$$

This implies

$$(t^{\frac{1}{2}} + t^{-\frac{1}{2}})V_K(t) \equiv (t^{\frac{1}{2}} + t^{-\frac{1}{2}})^r V_K(t)^r \pmod{(r, (t^{r+1} - 1)(t^{r-1} - 1))}.$$

Theorem 1.2(ii) is now proved. $\square$

1.4. EXAMPLES AND REMARKS.

We shall begin with the following proposition.

PROPOSITION 1.8. *For any knot K , either (i) or (ii) holds.*

$$(i) \quad V_K(t) - V_K(t^{-1}) \equiv 0 \pmod{(t^6 - 1)}.$$

$$(ii) \quad V_K(t) + V_K(t^{-1}) \equiv 0 \pmod{(t^2 - t + 1)}.$$

Proof. Write

$$V_K(t) = (t^2 - t + 1)Q(t) + at + b,$$

where $a, b \in Z$. Since $V_K(e^{\pi i/3})$ is a power of $i\sqrt{3}$ [13], it follows $a = 0$ or $a + 2b = 0$. If $a + 2b = 0$, a simple calculation shows (ii). If $a = 0$, it is clear that $t^2 - t + 1$ divides

$V_K(t) - V_K(t^{-1})$. Furthermore, $t^3 - 1$ and $t + 1$ also divide $V_K(t) - V_K(t^{-1})$ [13]. This proves (i). $\square$

By Proposition 1.8, it seems that Theorem 1.1 does not work in the case $r = 3$. However, we know Murasugi's result [29]

$$\Delta_K(t) \equiv (1 + t + \cdots + t^{\lambda-1})^{r-1} \Delta_k(t)^r \pmod{r},$$

where $\Delta_K(t)$ and $\Delta_k(t)$ denote Alexander polynomials of K and k respectively, and $\lambda = \text{lk}(K, \text{Fix } f)$. So we can evaluate $\text{lk}(K, \text{Fix } f)$ from Δ_K , and there is a possibility that Theorem 1.1 works in the case $r = 3$. But the author does not have such an example. In the cases $r \geq 5$, Theorem 1.1 works well. Here we give 2 examples.

Example. Consider $K = 10_{24}$ in [35]. Traczyk's criterion does not work for $r = 5$ because

$$V_K(t) - V_K(t^{-1}) \equiv 0 \pmod{(5, t^5 - 1)}.$$

However, we can apply Theorem 1.1 as follows. From $\Delta_K(t) \equiv (1 + t)^4 \pmod{5}$, if K has period 5, $\text{lk}(K, \text{Fix } f)$ must be 2. On the other hand, a simple calculation shows

$$V_K(t) + V_K(t^{-1}) \not\equiv 0 \pmod{(5, \frac{t^5 + 1}{t + 1})}.$$

Thus, by Theorem 1.1(ii), K can not have period 5.

Example. Consider $K = 10_{55}$ in [35]. Traczyk's criterion also does not work for $r = 5$. From $\Delta_K(t) \equiv 1 \pmod{5}$, if K has period 5, $\text{lk}(K, \text{Fix } f)$ must be 1. However, a simple calculation shows

$$V_K(t) - V_K(t^{-1}) \not\equiv 0 \pmod{(5, t^{10} - 1)}.$$

By Theorem 1.1(i), K can not have period 5.

Finally we remark that Theorem 1.2 holds for Jones polynomials of periodic links.

CHAPTER 2. SKEIN POLYNOMIALS OF PERIODIC KNOTS

The purpose of this Chapter is to present an unexpectedly strong regularity of "Skein polynomials" for knots with prime period, which was partially conjectured by P. Traczyk.

2.1. MAIN THEOREM.

Definition[8]. The *Skein polynomial* $P_L(v, z) \in Z[v^{\pm 1}, z^{\pm 1}]$ of an oriented link L is uniquely defined by

$$P_{\bigcirc}(v, z) = 1,$$

$$v^{-1}P \begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array} (v, z) - vP \begin{array}{c} \nwarrow \nearrow \\ \nearrow \searrow \end{array} (v, z) = zP \begin{array}{c} \longrightarrow \\ \longrightarrow \end{array} (v, z).$$

It is known that

$$\min\text{-deg}_z P_L(v, z) = 1 - \mu(L),$$

where $\mu(L)$ denotes the number of components of L [21]. So, we can decompose $P_L(v, z)$ as follows.

$$P_L(v, z) = \sum_{i \geq 0} P_i(v; L) z^{i+1-\mu(L)}.$$

Here, $P_i(v; L)$'s are in $Z[v^{\pm 1}]$. It is also known that $P_{2i+1}(v; L) = 0$ [21].

A pioneer work on this polynomial for periodic knots was demonstrated by P. Traczyk[40,41], which we now describe. He could notice an interesting property of a part of Skein polynomials and gave a handy, useful criterion for knots with prime period.

THEOREM 2.1[40,41]. *Let K be a knot with period r and k denote the linking number of K and the axis of rotation. Write*

$$(v^{-1} - v)P_0(v; K) = \sum_n \alpha_n v^n.$$

Then, $\alpha_n \equiv 0 \pmod{r}$ unless $n \equiv \pm k \pmod{r}$.

Example. Consider a knot K with period 7, made with 7 copies of a tangle depicted below. Then, we can evaluate

$$\begin{aligned} P_K(v, z) &= 7v^8 - 7v^{10} - 2v^{12} + 5v^{14} - 2v^{16} \\ &\quad + (14v^4 - 42v^6 + 77v^8 - 56v^{10} + 17v^{12} - 4v^{14} + 3v^{16})z^2 \\ &\quad + (49v^4 - 189v^6 + 301v^8 - 266v^{10} + 153v^{12} - 57v^{14} + 6v^{16})z^4 \\ &\quad + \dots \end{aligned}$$

Thus,

$$\begin{aligned}(v^{-1} - v)P_0(v; K) &= 7v^7 - 14v^9 + 5v^{11} + 7v^{13} - 7v^{15} + 2v^{17} \\ &\equiv 5v^{11} - 5v^{17} \pmod{7}.\end{aligned}$$

Note that $11 \equiv -3 \pmod{7}$ and $17 \equiv 3 \pmod{7}$.

P. Traczyk further conjectured that the regularity shown in Theorem 2.1 will also appear in higher terms $P_2(v; K)$, $P_4(v; K)$, ...

In this Chapter, we will give an answer for this and establish further regularity of Skein polynomials of periodic knots.

THEOREM 2.2. *If K is a knot with period r ,*

$$P_{2i}(v; K) \equiv p_{2i}P_0(v; K) \pmod{r}$$

for $2 \leq 2i \leq r - 3$, where p_{2i} 's are appropriate integers

We now check Theorem 2.2 with the example above. Then,

$$\begin{aligned}P_0(v; K) &= 7v^8 - 7v^{10} - 2v^{12} + 5v^{14} - 2v^{16} \\ &\equiv 5(v^{12} + v^{14} + v^{16}) \pmod{7},\end{aligned}$$

$$\begin{aligned}P_2(v; K) &= 14v^4 - 42v^6 + 77v^8 - 56v^{10} + 17v^{12} - 4v^{14} + 3v^{16} \\ &\equiv 3(v^{12} + v^{14} + v^{16}) \pmod{7},\end{aligned}$$

$$\begin{aligned}P_4(v; K) &= 49v^4 - 189v^6 + 301v^8 - 266v^{10} + 153v^{12} - 57v^{14} + 6v^{16} \\ &\equiv 6(v^{12} + v^{14} + v^{16}) \pmod{7}.\end{aligned}$$

Thus, we have $P_2(v; K) \equiv 2P_0(v; K) \pmod{7}$ and $P_4(v; K) \equiv 4P_0(v; K) \pmod{7}$.

To prove Theorem 2.2, we have to investigate Skein polynomials of torus knots and links which are typical knots and links with periods. We shall start by introducing "Hecke algebras", which will help our work. In sections 2.3 and 2.4, as an essential part of the proof, we will compute Skein polynomials of torus knots and links. Section 2.5 will be then devoted to the proof of Theorem 2.2 using results obtained in sections 2.2 and 2.3.

2.2. HECKE ALGEBRAS.

We will first introduce "Hecke algebras". As in the case of Jones', Skein polynomials can be described as a weighted sum of irreducible characters of Hecke algebras.

Definition. Suppose n is a positive integer and $q \neq 0$ is not a root of unity. Then, Hecke algebra $H_n(q)$ will be defined as an algebra over $\mathbb{C}$ with generators

$$1, g_1, g_2, \dots, g_{n-1}$$

and relations

$$g_i g_j = g_j g_i \quad \text{if } |i - j| > 1,$$

$$g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1},$$

$$g_i - g_i^{-1} = (q - q^{-1}).$$

As is well-known, $H_n(q)$ is semisimple. Furthermore, simple components of $H_n(q)$ are indexed by "partitions" of n . A partition can be represented as follows.

Such a diagram will be called *Young diagram* if the length of rows in the diagram made to be nonincreasing. We denote the set of Young diagrams with n nodes by Λ_n .

PROPOSITION 2.3[43]. $H_n(q)$ decomposes into a direct sum of full matrix algebras $H_{n,\lambda}$, where $\lambda \in \Lambda_n$.

By $\rho_{n,\lambda}$, we shall denote an irreducible representation of $H_n(q)$ corresponding to λ . Then, we can evaluate the dimension of $\rho_{n,\lambda}$ as follows.

Each box of a Young diagram defines a *hook* consisting of boxes horizontally to the right and vertically below. The number of boxes in a hook will be called *hook length*.

PROPOSITION 2.4[43]. The dimension $d_{n,\lambda}$ of $\rho_{n,\lambda}$ is $n!$ divided by the product of hook lengths of λ .

Another view for $d_{n,\lambda}$ will be possible. For $\lambda, \mu \in \Lambda_n$, a sequence of Young diagrams

$$\lambda = \nu_0, \nu_1, \dots, \nu_f = \mu$$

is called a *path* of order f from λ to μ if every ν_{i+1} is obtained from ν_i by removing or adding one node. Then, $d_{n,\lambda}$ can be also evaluated as the number of paths of order n from λ to $\phi \in \Lambda_0$ [43].

We can explicitly write down representation matrices for $(g_1 g_2 \dots g_{n-1})^n$. In fact, we can observe this element is in the center of $H_n(q)$ and so, $\rho_{n,\lambda}(g_1 g_2 \dots g_{n-1})^n$'s are scalars given as follows.

PROPOSITION 2.5[13]. By $u_{n,\lambda}$ and $v_{n,\lambda}$, we denote the numbers of paths of order $n-2$ from λ to $\square\square$ and to $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$ respectively. Then, we have

$$\rho_{n,\lambda}(g_1 g_2 \dots g_{n-1})^n = q^{n(n-1)(u_{n,\lambda} - v_{n,\lambda})/d_{n,\lambda}}.$$

This will play an important role in the next section. We can evaluate $u_{n,\lambda}$ and $v_{n,\lambda}$ for certain λ . Let Λ'_n be the set of Young diagrams with the following shape.

PROPOSITION 2.6. For $\lambda \in \Lambda'_n$ above,

$$u_{n,\lambda} = \binom{\beta + \gamma - 1}{\gamma}, \quad v_{n,\lambda} = \binom{\beta + \gamma - 1}{\beta}.$$

So, we have

$$d_{n,\lambda} = u_{n,\lambda} + v_{n,\lambda} = \binom{\beta + \gamma}{\gamma}.$$

2.3. TORUS LINKS.

To perform computations for torus links, we shall recall the character formula for Skein polynomials given in [13].

Let B_n denote the n -string braid group with standard generators

$$\sigma_1, \sigma_2, \dots, \sigma_{n-1}.$$

For a Young diagram λ with n nodes, we also denote by $\rho_{n,\lambda}$ an irreducible representation of B_n which arises from $\rho_{n,\lambda}$ with an obvious homomorphism $\sigma_i \rightarrow g_i$. Characters of $\rho_{n,\lambda}$'s, for simplicity, will be denoted by $\chi_{n,\lambda}$'s.

Now, we define $R_\lambda(v, q)$ as the product of terms covered by λ on the following diagram.

$$\begin{array}{cccc} v^{-1} - v & v^{-1}q - vq^{-1} & v^{-1}q^2 - vq^{-2} & \dots \\ v^{-1}q^{-1} - vq & v^{-1} - v & v^{-1}q - vq^{-1} & \dots \\ v^{-1}q^{-2} - vq^2 & v^{-1}q^{-1} - vq & v^{-1} - v & \dots \\ \dots & \dots & \dots & \dots \end{array}$$

Furthermore, we fill in each box with hook length in λ , and replace each integer appearing, say m , by $q^m - q^{-m}$. Then $Q_\lambda(q)$ is defined as the product of these terms. For instance, suppose λ is of the following shape.

Then, $R_\lambda(v, q)$ becomes

$$(v^{-1} - v)(v^{-1}q - vq^{-1})(v^{-1}q^2 - vq^{-2})(v^{-1}q^{-1} - vq)(v^{-1} - v)(v^{-1}q^{-2} - vq^2)$$

and $Q_\lambda(q)$ becomes

$$(q^5 - q^{-5})(q^3 - q^{-3})(q - q^{-1})(q^3 - q^{-3})(q - q^{-1})(q - q^{-1}).$$

For simplicity, we denote $q^m - q^{-m}$ and $\frac{q^m - q^{-m}}{q - q^{-1}}$ by $[m]$ and $[\bar{m}]$ respectively.

PROPOSITION 2.7[13]. For $\xi \in B_n$, the closure $\hat{\xi}$ represents an oriented link L in S^3 . Let $\epsilon(\xi)$ denote the exponent sum of ξ . Then,

$$\frac{v^{-1} - v}{[1]} P_L(v, [1]) = v^{\epsilon(\xi)} \sum_{\lambda \in \Lambda_n} \frac{R_\lambda(v, q)}{Q_\lambda(q)} \chi_{n,\lambda}(\xi).$$

We now consider the (r, nr) torus link L_n . As L_n is represented by the closure of $(\sigma_1 \sigma_2 \dots \sigma_{r-1})^{nr} \in B_r$, it follows from Propositions 2.5 and 2.7 that

PROPOSITION 2.8.

$$P_{L_n}(v, [1]) = \frac{[1]}{v^{-1} - v} \sum_{\lambda} \frac{R_{\lambda}(v, q)}{Q_{\lambda}(q)} d_{n, \lambda} q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}}.$$

Using this, we will evaluate Skein polynomials of torus links. We begin with the following.

PROPOSITION 2.9.

$$z^{r-3}(P_{L_n}(v, z) - v^{nr(r-1)}P_{L_0}(v, z)) = \sum_{i \geq 1} P_{2i}(v; L_n) z^{2i-2}.$$

Proof. Note that L_0 is a trivial link with r components and so,

$$P_{L_0}(v, z) = \left(\frac{v^{-1} - v}{z}\right)^{r-1}.$$

On the other hand, we have

$$P_{L_n}(v, z) = v^{nr(r-1)} \left(\frac{v^{-1} - v}{z}\right)^{r-1} + \sum_{i \geq 1} P_{2i}(v; L_n) z^{2i+1-r}$$

[21]. Thus, required equality is now apparent. $\square$

We must evaluate the left hand side in another way.

PROPOSITION 2.10. $[1]^{r-3}(P_{L_n}(v, [1]) - v^{nr(r-1)}P_{L_0}(v, [1]))$ should be a polynomial in $Z[v^{\pm 1}, q^{\pm 1}]$ of the form

$$[\bar{r}] \frac{R_1(v, q)}{Q_1(q)} + r \frac{R_2(v, q)}{Q_2(q)},$$

where $R_1, R_2 \in Z[v^{\pm 1}, q^{\pm 1}]$ and $Q_1, Q_2 \in Z[q^{\pm 1}]$. Furthermore, neither $Q_1(1)$ nor $Q_2(1)$ is divisible by r .

Proof. By Proposition 2.8, we have

$$\begin{aligned} & [1]^{r-3}(P_{L_n}(v, [1]) - v^{nr(r-1)}P_{L_0}(v, [1])) \\ &= v^{nr(r-1)} \frac{[1]^{r-2}}{v^{-1} - v} \sum_{\lambda} \frac{R_{\lambda}(v, q)}{Q_{\lambda}(q)} d_{r, \lambda} (q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}} - 1). \end{aligned}$$

We shall divide the summation on the right hand side into 2 parts. The proof will be completed by 2 claims below. $\square$

CLAIM 2.10.1. The first term

$$[\bar{r}] \frac{R_1(v, q)}{Q_1(q)}$$

in Proposition 2.10 should be

$$v^{nr(r-1)} \frac{[1]^{r-2}}{v^{-1} - v} \sum_{\lambda \in \Lambda'_r} \frac{R_\lambda(v, q)}{Q_\lambda(q)} d_{r, \lambda} (q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}} - 1).$$

Proof of Claim 2.10.1. Suppose $\lambda, \lambda^* \in \Lambda'_r$ have the following shapes.

If $\gamma = \beta$, we have $u_{r, \lambda} = v_{r, \lambda}$ and

$$q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}} - 1 = 0.$$

So, we suppose $\gamma > \beta$. A direct calculation shows

$$R_\lambda(v, q) = \prod_{i=-\gamma}^{\beta} (v^{-1} q^i - v q^i), \quad Q_\lambda(q) = [\beta]! [\gamma]! [r],$$

where $[m]!$ stands for $[1][2] \dots [m]$. Similarly, we have

$$R_{\lambda^*}(v, q) = \prod_{i=-\beta}^{\gamma} (v^{-1} q^i - v q^i), \quad Q_{\lambda^*}(q) = [\beta]! [\gamma]! [r].$$

Furthermore, by Proposition 2.6,

$$d_{r, \lambda} = \binom{\beta + \gamma}{\gamma} = \binom{\beta + \gamma}{\beta} = d_{r, \lambda^*}$$

and

$$\begin{aligned} u_{r, \lambda} - v_{r, \lambda} &= \binom{\beta + \gamma - 1}{\gamma} - \binom{\beta + \gamma - 1}{\beta} \\ &= \frac{(\beta - \gamma)}{\beta + \gamma} d_{r, \lambda} \\ &= -\frac{(\gamma - \beta)}{\beta + \gamma} d_{r, \lambda^*} \\ &= -(u_{r, \lambda^*} - v_{r, \lambda^*}). \end{aligned}$$

Therefore, in the summation of Claim 2.10.1, the contribution of λ, λ^* is

$$\begin{aligned} & \frac{R_\lambda(v, q)}{Q_\lambda(q)} d_{r, \lambda} (q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}} - 1) \\ & + \frac{R_{\lambda^*}(v, q)}{Q_{\lambda^*}(q)} d_{r, \lambda^*} (q^{nr(r-1)(u_{r, \lambda^*} - v_{r, \lambda^*})/d_{r, \lambda^*}} - 1) \\ & = \frac{\binom{\beta+\gamma}{\gamma} (R_\lambda(v, q)(q^{(\beta-\gamma)nr} - 1) + R_{\lambda^*}(v, q)(q^{(\gamma-\beta)nr} - 1))}{[\beta]![\gamma]![r]} \end{aligned}$$

and this numerator becomes

$$(q^{(\beta-\gamma)nr} - 1) \prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i}) + (q^{(\gamma-\beta)nr} - 1) \prod_{i=-\beta}^{\gamma} (v^{-1}q^i - vq^{-i}).$$

As r is odd, $\beta - \gamma$ is even and so, $q^{(\beta-\gamma)nr} - 1$ and $q^{(\gamma-\beta)nr} - 1$ are always divisible by $[r]$. Hence, to prove Claim 2.10.1, it suffices to show

$$\prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i}) \equiv \prod_{i=-\beta}^{\gamma} (v^{-1}q^i - vq^{-i}) \pmod{[r]}.$$

In fact, the left hand side can be modified as

$$\begin{aligned} & \prod_{i=-\beta}^{\beta} (v^{-1}q^i - vq^{-i}) q^{(\gamma-\beta)r} \prod_{i=-\gamma}^{-\beta-1} (v^{-1}q^i - vq^{-i}) \\ & \equiv \prod_{i=-\beta}^{\beta} (v^{-1}q^i - vq^{-i}) \prod_{i=-\gamma}^{-\beta-1} (v^{-1}q^{r+i} - vq^{-r-i}) \\ & \equiv \prod_{i=-\beta}^{\beta} (v^{-1}q^i - vq^{-i}) \prod_{i=\beta+1}^{\gamma} (v^{-1}q^i - vq^{-i}) \\ & \equiv \prod_{i=-\beta}^{\gamma} (v^{-1}q^i - vq^{-i}) \pmod{[r]}. \end{aligned}$$

Thus, we have done. $\square$

CLAIM 2.10.2. The second term

$$r \frac{R_2(v, q)}{Q_2(q)}$$

in Proposition 2.10 should be

$$v^{nr(r-1)} \frac{[1]^{r-2}}{v^{-1} - v} \sum_{\lambda \notin \Lambda'_r} \frac{R_\lambda(v, q)}{Q_\lambda(q)} d_{r, \lambda} (q^{nr(r-1)(u_{r, \lambda} - v_{r, \lambda})/d_{r, \lambda}} - 1).$$

Proof of Claim 2.10.2. If $\lambda \notin \Lambda'_r$, r can not appear as a hook length in λ . So, $Q_\lambda(q)$ can not have factor $[r]$ and, if we write, $Q_2(q)$ can not have the factor $[\bar{r}]$. Furthermore, from the hook length formula, $d_{r,\lambda}$ should be divisible by r . Claim 2.10.2 now follows immediately. $\square$

We shall close this section with the following Lemma.

LEMMA 2.11. For $2 \leq 2i \leq r-1$,

$$P_{2i}(v; L_n) \equiv 0 \pmod{r}.$$

Proof. By Propositions 2.9 and 2.10, we have

$$\sum_{i \geq 1} P_{2i}(v; L_n) [1]^{2i-2} = [\bar{r}] \frac{R_1(v, q)}{Q_1(q)} + r \frac{R_2(v, q)}{Q_2(q)}.$$

Derivate both sides with respect to q for $2i$ times, where $0 \leq 2i \leq r-3$. It should be noted that

$$\lim_{q \rightarrow 1} \frac{d^m}{dq^m} [\bar{r}] \equiv 0 \pmod{r}$$

for $0 \leq m \leq r-2$ and this is false if $m \geq r-1$. Therefore, by taking a limit $q \rightarrow 1$, the left hand side will turn into a linear combination of

$$P_2(v; L_n), P_4(v; L_n), \dots, P_{2i+2}(v; L_n)$$

and the right hand side will vanish mod r . Thus, we can prove Lemma 2.11 inductively. $\square$

2.4. TORUS KNOTS.

In this section, we will detect Skein polynomials of torus knots. Recall Jones' result, which can be derived from Proposition 2.7 and representations of braid groups. See also Chapter 3.

THEOREM 2.12[13]. Let K be the (m, n) torus knot, where we suppose m and n are relatively prime and n is positive. Then, we have

$$\frac{v^{-1} - v}{[1]} P_K(v, [1]) = v^{m(n-1)} \Sigma_v(m, n),$$

where

$$\Sigma_v(m, n) = \sum_{\beta + \gamma + 1 = n} (-1)^\gamma q^{(\beta - \gamma)m} \frac{\prod_{i=-\gamma}^{\beta} (v^{-1} q^i - v q^{-i})}{[\beta]! [\gamma]! [n]}.$$

We shall prepare an interesting identity here.

PROPOSITION 2.13.

$$\Sigma_v(0, n) = \frac{v^{-n} - v^n}{[n]}.$$

Proof. Note that

$$\Sigma_v(\pm 1, n) = v^{\mp(n-1)} \frac{v^{-1} - v}{[1]}$$

by Theorem 2.12 because $(\pm 1, n)$ torus knots are trivial. We can directly check Proposition 2.13 in cases $n = 1, 2$. If $n > 2$,

$$\begin{aligned} & [n]((q^{n-1} + q^{-n+1})\Sigma_v(0, n) - \Sigma_v(1, n) - \Sigma_v(-1, n)) \\ &= \sum_{\beta+\gamma+1=n} (-1)^\gamma (q^{n-1} + q^{-n+1} - q^{\beta-\gamma} - q^{\gamma-\beta}) \frac{\prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i})}{[\beta]![\gamma]!} \\ &= \sum_{\beta+\gamma+1=n} (-1)^\gamma [\beta][\gamma] \frac{\prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i})}{[\beta]![\gamma]!} \\ &= - \sum_{\beta+\gamma+1=n, \beta, \gamma \geq 1} (-1)^{\gamma-1} (v^{-1}q^{-\gamma} - vq^\gamma)(v^{-1}q^\beta - vq^{-\beta}) \frac{\prod_{i=-\gamma+1}^{\beta-1} (v^{-1}q^i - vq^{-i})}{[\beta-1]![\gamma-1]!} \\ &= - \sum_{\beta'+\gamma'+1=n-2} (-1)^{\gamma'} (v^{-2}q^{\beta'-\gamma'} + v^2q^{\gamma'-\beta'} - q^{n-1} - q^{-n+1}) \frac{\prod_{i=-\gamma'}^{\beta'} (v^{-1}q^i - vq^{-i})}{[\beta']![\gamma']!} \\ &= [n-2](q^{n-1} + q^{-n+1})\Sigma_v(0, n-2) - [n-2](v^{-2}\Sigma_v(1, n-2) + v^2\Sigma_v(-1, n-2)). \end{aligned}$$

By the hypothesis of induction, the first term should be

$$[n-2](q^{n-1} + q^{-n+1}) \frac{v^{n-2} - v^{-n+2}}{[n-2]} = (q^{n-1} + q^{-n+1})(v^{n-2} - v^{-n+2}).$$

Furthermore,

$$\begin{aligned} & [n]\Sigma_v(\pm 1, n) - v^{\mp 2}[n-2]\Sigma_v(\pm 1, n-2) \\ &= [n]v^{\mp(n-1)} \frac{(v^{-1} - v)}{[1]} - [n-2]v^{\mp 2}v^{\mp(n-3)} \frac{(v^{-1} - v)}{[1]} \\ &= (q^{n-1} + q^{-n+1})v^{\mp(n-1)}(v^{-1} - v). \end{aligned}$$

Thus, we have

$$[n](q^{n-1} + q^{-n+1})\Sigma_v(0, n) = (q^{n-1} + q^{-n+1})(v^{-n} - v^n).$$

That is,

$$\Sigma_v(0, n) = \frac{v^{-n} - v^n}{[n]}. \quad \square$$

Let T_n denote the $(r, nr+k)$ torus knot, where $1 \leq k \leq r-1$. We begin with the case $n = 0$ as a special one.

PROPOSITION 2.14.

$$q^{(k-1)r}[\bar{k}]P_{T_0}(v, [1]) \equiv v^{(k-1)r} \frac{v^{-k} - v^k}{v^{-1} - v} \pmod{[\bar{r}]}.$$

Proof. By Theorem 2.12 and Proposition 2.13, we have

$$\begin{aligned} & q^{(k-1)r}[\bar{k}]P_{T_0}(v, [1]) - v^{(k-1)r} \frac{v^{-k} - v^k}{v^{-1} - v} \\ &= v^{(k-1)r} \frac{1}{v^{-1} - v} \sum_{\beta+\gamma+1=k} (-1)^\gamma (q^{(k-1)r+(\beta-\gamma)r} - 1) \frac{\prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i})}{[\beta]![\gamma]!} \\ &= v^{(k-1)r} \frac{1}{v^{-1} - v} \sum_{\beta+\gamma+1=k} (-1)^\gamma (q^{2\beta r} - 1) \frac{\prod_{i=-\gamma}^{\beta} (v^{-1}q^i - vq^{-i})}{[\beta]![\gamma]!}. \end{aligned}$$

Obviously, the left hand side is a polynomial in $Z[v^{\pm 1}, q^{\pm 1}]$. Analyzing the right hand side, we can show this is divisible by $[\bar{r}]$. In fact, each $q^{2\beta r} - 1$ is divisible by $[r]$ so that $[\bar{r}]$ should be a common factor for all terms. This completes the proof of Proposition 2.14. $\square$

PROPOSITION 2.15. For $2 \leq 2i \leq r-3$, we have

$$P_{2i}(v; T_0) \equiv p_{2i}P_0(v; T_0) \pmod{r},$$

where p_{2i} 's are appropriate integers.

Proof. By Proposition 2.14, we have

$$q^{(k-1)r}[\bar{k}]P_{T_0}(v, [1]) \equiv v^{r(k-1)} \frac{v^{-k} - v^k}{v^{-1} - v} \pmod{[\bar{r}]}.$$

Derivate both sides with respect to q for $2i$ times, where $0 \leq 2i \leq r-3$. By taking a limit $q \rightarrow 1$, the left hand side will turn into a linear combination of

$$P_0(v; T_0), P_2(v; T_0), \dots, P_{2i}(v; T_0)$$

and the right hand side will vanish mod r as in Lemma 2.11. Thus, we can prove Proposition 2.15 inductively. $\square$

Let us consider $n = -1$ as another special case.

PROPOSITION 2.16. For $2 \leq 2i \leq r-3$, we have

$$P_{2i}(v; T_{-1}) \equiv p_{2i}P_0(v; T_{-1}) \pmod{r},$$

where p_{2i} 's are the same integers as in Proposition 2.15.

Before the proof, we should note

PROPOSITION 2.17. Let T_n^* denote the $(r, nr - k)$ torus knot. Then,

$$P_{2i}(v; T_n^*) = P_{2i}(v^{-1}; T_{-n}).$$

Proof. Since T_n^* is the mirror image of T_{-n} , we have

$$P_{T_n^*}(v, z) = P_{T_{-n}}(v^{-1}, -z)$$

[21]. Now, Proposition 2.17 follows immediately. $\square$

Proof of Proposition 2.16. By Proposition 2.17, it suffices to show for T_1^* . By Proposition 2.15, we have

$$q^{(r-k-1)r} [r-k] P_{T_1^*}(v, [1]) \equiv v^{(r-k-1)r} \frac{v^{-r+k} - v^{r-k}}{v^{-1} - v} \pmod{[r]}.$$

Here, a simple calculation shows that

$$q^{(r-k-1)r} [r-k] \equiv -q^{kr} [k] \pmod{[r]}$$

and so, we have

$$-q^{kr} [\bar{k}] P_{T_1^*}(v, [1]) \equiv v^{(r-k-1)r} \frac{v^{-r+k} - v^{r-k}}{v^{-1} - v} \pmod{[\bar{r}]}.$$

As in Proposition 2.15, derivate the left hand side with respect to q for $2i$ times, where $0 \leq 2i \leq r-3$. Taking a limit $q \rightarrow 1$, we will also have a linear combination of

$$P_0(v; T_1^*), P_2(v; T_1^*), \dots, P_{2i}(v; T_1^*)$$

with coefficients equal to ones in Proposition 2.15 $\pmod{r}$. Thus, we can deduce Proposition 2.16 inductively. $\square$

Furthermore, we have

PROPOSITION 2.18. The polynomial

$$v^{-r(r-1)} P_{T_{n+1}}(v, [1]) - 2P_{T_n}(v, [1]) + v^{r(r-1)} P_{T_{n-1}}(v, [1])$$

is divisible by $[\bar{r}]$. Hence, for $0 \leq 2i \leq r-3$, it follows

$$v^{-r(r-1)} P_{2i}(v; T_{n+1}) - 2P_{2i}(v; T_n) + v^{r(r-1)} P_{2i}(v; T_{n-1}) \equiv 0 \pmod{r}.$$

Proof. By Theorem 2.12, we have

$$P_{T_n}(v, [1]) = v^{(r-1)(nr+k)} \frac{[1]}{v^{-1} - v} \sum_{\beta+\gamma+1=r} (-1)^\gamma q^{(\beta-\gamma)(nr+k)} \frac{\prod_{i=-\gamma}^{\beta} (v^{-1} q^i - v q^{-i})}{[\beta]! [\gamma]! [r]}.$$

Hence, it follows that

$$\begin{aligned}
& v^{-r(r-1)} P_{T_{n+1}}(v, [1]) - 2P_{T_n}(v, [1]) + v^{r(r-1)} P_{T_{n-1}}(v, [1]) \\
&= v^{(r-1)(nr+k)} \frac{[1]}{v^{-1} - v} \\
&\quad \times \sum_{\beta+\gamma+1=r} (-1)^\gamma (q^{(\beta-\gamma)r} - 2 + q^{(\gamma-\beta)r}) q^{\beta(nr+k)} \frac{\prod_{i=-\gamma}^{\beta} (v^{-1} q^i - v q^{-i})}{[\beta]! [\gamma]! [r]} \\
&\equiv 0 \pmod{[r]}
\end{aligned}$$

because each $q^{(\beta-\gamma)r} - 2 + q^{(\gamma-\beta)r}$ is divisible by $[r]^2$. Now, Proposition 2.18 follows immediately. $\square$

From Propositions 2.15, 2.16 and 2.18, we can deduce

LEMMA 2.19. For $0 \leq 2i \leq r-3$, we have

$$P_{2i}(v; T_n) \equiv p_{2i} P_0(v; T_n) \pmod{r},$$

where p_{2i} 's are the same integers as in Proposition 2.15.

As a direct consequence of Proposition 2.17 and Lemma 2.19, we also have

LEMMA 2.20. For $0 \leq 2i \leq r-3$, we have

$$P_{2i}(v; T_n^*) \equiv p_{2i} P_0(v; T_n^*) \pmod{r},$$

where p_{2i} 's are the same integers as in Proposition 2.15.

2.5. PERIODIC REDUCTIONS.

The arguments used here are based on Traczyk[40] and so we follow the proofs briefly. For details, see [40].

LEMMA 2.21. Let L be a link with r components, having period r with no component invariant under rotation. Then, for $2 \leq 2i \leq r-1$,

$$P_{2i}(v; L) \equiv 0 \pmod{r}.$$

Proof. L has a diagram D with period r in a plane. Let $D_{+...+}$, $D_{-...-}$ and $D_{0,...,0}$ denote diagrams obtained from D by making all the crossings in a periodic orbit positive, negative and nullified respectively. We will call an operation $D_{+...+} \rightarrow D_{-...-}$ or its inverse a *periodic reduction*. We now use a congruence[34]

$$v^{-r} P_{D_{+...+}}(v, z) - v^r P_{D_{-...-}}(v, z) \equiv z^r P_{D_{0,...,0}}(v, z) \pmod{r}.$$

For example, in the case $r = 3$, we have the following decomposition for Skein polynomials.

$$\begin{array}{ccccccc}
 & & & & D_{+++} & & \\
 & & & & & & \\
 & & D_{-++} & & D_{0++} & & \\
 & D_{--+} & D_{-0+} & D_{0-+} & D_{00+} & & \\
 D_{---} & D_{--0} & D_{-0-} & D_{-00} & D_{0--} & D_{0-0} & D_{00-} & D_{000}
 \end{array}$$

Using symmetries, it follows

$$\begin{aligned}
 P_{D_{+++}}(v, z) &= v^6 P_{D_{---}}(v, z) + 3v^5 z P_{D_{0--}}(v, z) \\
 &\quad + 3v^4 z^2 P_{D_{-00}}(v, z) + v^3 z^3 P_{D_{000}}(v, z) \\
 &\equiv v^6 P_{D_{---}}(v, z) + v^3 z^3 P_{D_{000}}(v, z) \pmod{3}.
 \end{aligned}$$

The number of components in $D_{0\dots 0}$ should be 2 or $2r$ from a combinatorics. Recall

$$\min\text{-deg}_z P_L(v, z) = 1 - \mu(L)$$

for a link L . So, if $D_{0\dots 0}$ has 2 components, the congruence above tells us

$$v^{-r} P_{2i}(v; D_{+,\dots,+}) - v^r P_{2i}(v; D_{-,\dots,-}) \equiv 0 \pmod{r}$$

for $0 \leq 2i < 2(r-1)$. If $D_{0\dots 0}$ has $2r$ components, then

$$v^{-r} P_{2i}(v; D_{+,\dots,+}) - v^r P_{2i}(v; D_{-,\dots,-}) \equiv P_{2i}(v; D_{0\dots 0}) \pmod{r}.$$

We will first show Lemma 2.21 in the following case.

Case 2.21.1. D is a closed nr braid.

We prove Lemma 2.21 by induction on n in this case. Suppose D is represented as the closure of $\xi^r \in B_{nr}$, where $\xi \in B_{nr}$. From the assumption for D , ξ must represent an nr cycle if we regard $\xi \in B_{nr}$ as an element of symmetric group S_{nr} naturally. Therefore, a sequence of periodic reductions reduces D into the (r, nr) torus link [40] and so, to prove Lemma 2.21 in this case, it suffices to show Lemma 2.21 holds for $D_{+,\dots,+}$ if and only if it does for $D_{-,\dots,-}$ by virtue of Lemma 2.11.

If $D_{0\dots 0}$ has 2 components, there is nothing to do. We suppose $D_{0\dots 0}$ has $2r$ components. To prove our assertion, it is sufficient to show

$$P_{2i}(v; D_{0\dots 0}) \equiv 0 \pmod{r}$$

for $0 \leq 2i \leq r-1$. It should be noted that $D_{0\dots 0}$ consists of 2 link diagrams, say $D'_{0\dots 0}, D''_{0\dots 0}$, satisfying the assumption in Lemma 2.21. Let $D_{0\dots 0, +,\dots,+}, D_{0\dots 0, -,\dots,-}$ and $D_{0\dots 0, 0\dots 0}$ denote diagrams obtained from $D_{0\dots 0}$ by making all crossings in a periodic

orbit of a crossing between $D'_{0\dots 0}$ and $D''_{0\dots 0}$ positive, negative and nullified respectively. Then, the number of components of $D_{0\dots 0,0\dots 0}$ must be r from a combinatorics and so,

$$v^{-r}P_{2i}(v; D_{0\dots 0,+ \dots +}) - v^r P_{2i}(v; D_{0\dots 0,- \dots -}) \equiv 0 \pmod{r}$$

for $0 \leq 2i < 2r$. Repeating this operation, we can make $D_{0\dots 0}$ into a disjoint union $D'_{0\dots 0} \cup D''_{0\dots 0}$. This implies that

$$P_{2i}(v; D_{0\dots 0}) \equiv v^{2\nu r} P_{2i}(v; D'_{0\dots 0} \cup D''_{0\dots 0}) \pmod{r}$$

for $0 \leq 2i < 2r$, where νr equals the linking number of $D'_{0\dots 0}$ and $D''_{0\dots 0}$. Since

$$P_{D'_{0\dots 0} \cup D''_{0\dots 0}}(v, z) = \frac{v^{-1} - v}{z} P_{D'_{0\dots 0}}(v, z) P_{D''_{0\dots 0}}(v, z)$$

[21], we have

$$P_{2i}(v; D_{0\dots 0}) \equiv v^{2\nu r} (v^{-1} - v) \sum_{j=0}^i P_{2j}(v; D'_{0\dots 0}) P_{2i-2j}(v; D''_{0\dots 0}) \pmod{r}.$$

This holds for $0 \leq 2i < 2r$, however we need only for $0 \leq 2i \leq r-1$. Note that both of $D'_{0\dots 0}$ and $D''_{0\dots 0}$ are closed braids with strings less than nr . By the hypothesis of induction, the right hand side vanishes for $2 \leq 2i \leq r-1$ and our assertion is true.

We will complete the proof of Lemma 2.21 by showing for the case D is not a closed braid.

Case 2.21.2. Otherwise.

If D has no crossing point, then

$$P_D(v, z) = \left(\frac{v^{-1} - v}{z} \right)^{r-1}$$

and hence Lemma 2.21 clearly holds. Now, as above, induction on the number of crossings in D will reduce this case into previous case and then, the proof of Lemma 2.21 will be completed. $\square$

Proof of Theorem 2.2. Suppose K is a knot with period r and k , $1 \leq k \leq r-1$, denotes mod r linking number of K and the axis of rotation. K has a diagram with period r in a plane. Define $D_{+ \dots +}$, $D_{- \dots -}$ and $D_{0\dots 0}$ as in previous Lemma. We can now observe the number of components of $D_{0\dots 0}$ should be 2 or $r+1$.

If $D_{0\dots 0}$ has 2 components, then

$$v^{-r}P_{2i}(v; D_{+ \dots +}) - v^r P_{2i}(v; D_{- \dots -}) \equiv 0 \pmod{r}$$

for $0 \leq 2i \leq r-3$ as before. If $D_{0\dots 0}$ has $r+1$ components, then it too follows

$$v^{-r}P_{2i}(v; D_{+ \dots +}) - v^r P_{2i}(v; D_{- \dots -}) \equiv P_{2i}(v; D_{0\dots 0}) \pmod{r}$$

as before.

Case 2.2.1. D is a closed $nr \pm k$ braid.

As in Lemma 2.21, we can also prove Theorem 2.2 by induction on n in this case. As was observed, a sequence of periodic reductions will reduce D into a diagram of the $(r, nr \pm k)$ torus knot [40]. Thus, it suffices to show that Theorem 2.2 holds for $D_{+...+}$ if and only if it does for $D_{-...-}$ by virtue of Lemmas 2.19 and 2.20.

If $D_{0...0}$ has 2 components, this clearly holds from the congruence above. We suppose $D_{0...0}$ has $r+1$ components. Note that $D_{0...0}$ consists of a diagram $D'_{0...0}$ of a knot with period r and a diagram $D''_{0...0}$ of a link satisfying the assumption in Lemma 2.21. We can prove

$$P_{2i}(v; D_{0...0}) \equiv v^{2\nu r} P_{2i}(v; D'_{0...0} \cup D''_{0...0}) \pmod{r}$$

for $0 \leq 2i < 2r$ as in the proof of Lemma 2.21, and we have

$$\begin{aligned} P_{2i}(v; D'_{0...0} \cup D''_{0...0}) &= (v^{-1} - v) \sum_{j=0}^i P_{2j}(v; D'_{0...0}) P_{2i-2j}(v; D''_{0...0}) \\ &\equiv (v^{-1} - v) P_0(v; D''_{0...0}) P_{2i}(v; D'_{0...0}) \pmod{r} \end{aligned}$$

for $0 \leq 2i \leq r-1$ by using Lemma 2.21, where νr equals the linking number of $D'_{0...0}$ and $D''_{0...0}$. Note that $D'_{0...0}$ is also a closed braid with strings less than $nr \pm k$. Furthermore, $\pmod{r}$ linking number must be preserved even after $D''_{0...0}$ is removed. By the hypothesis of induction, for $2 \leq 2i \leq r-3$, we have

$$\begin{aligned} &P_{2i}(v; D_{0...0}) - p_{2i} P_0(v; D_{0...0}) \\ &\equiv v^{2\nu r} (P_{2i}(v; D'_{0...0} \cup D''_{0...0}) - p_{2i} P_0(v; D'_{0...0} \cup D''_{0...0})) \\ &\equiv v^{2\nu r} (v^{-1} - v) P_0(v; D''_{0...0}) (P_{2i}(v; D'_{0...0}) - p_{2i} P_0(v; D'_{0...0})) \\ &\equiv 0 \pmod{r}. \end{aligned}$$

Equivalently,

$$v^{-r} (P_{2i}(v; D_{+...+}) - p_{2i} P_0(v; D_{+...+})) \equiv v^r (P_{2i}(v; D_{-...-}) - p_{2i} P_0(v; D_{-...-})) \pmod{r}.$$

This guarantees our assertion.

Case 2.2.1 Otherwise.

As in Lemma 2.21, induction on the number of crossings will reduce this case into previous case. We can then complete the proof of Theorem 2.2. $\square$

CHAPTER 3. KAUFFMAN POLYNOMIALS OF PERIODIC KNOTS

In this Chapter, we shall discuss another regularity appearing in “Kauffman polynomials” of periodic knots. Surprisingly, we will discover a quite strong one analogous to Skein’s.

3.1. MAIN THEOREM.

We now introduce “Kauffman polynomials” which is known as another generalization of Jones.

Definition[17]. Let D be a diagram of an oriented link L with μ components. The algebraic crossing number of D will be denoted by $w(D)$. Then, the *Kauffman polynomial* $F_L(a, z) \in Z[a^{\pm 1}, z^{\pm 1}]$ of L is defined by

$$a^{-w(D)} \Lambda_D(a, z),$$

where $\Lambda_D(a, z)$ is a regular isotopy invariant of diagrams of L with properties

$$\begin{aligned} \Lambda_{\bigcirc}(a, z) &= 1, \\ \Lambda_{\text{positive crossing}}(a, z) &= a \Lambda_{\text{negative crossing}}(a, z), \quad \Lambda_{\text{negative crossing}}(a, z) = a^{-1} \Lambda_{\text{positive crossing}}(a, z), \\ \Lambda_{\text{crossing}}(a, z) - \Lambda_{\text{other crossing}}(a, z) &= z(\Lambda_{\text{smooth}}(a, z) - \Lambda_{\text{other smooth}}(a, z)). \end{aligned}$$

It is known that

$$\min\text{-deg}_z F_L(a, z) = 1 - \mu(L),$$

where $\mu(L)$ denotes the number of components in L [17]. So, as in Skein polynomials, we shall decompose $F_L(a, z)$ as follows.

$$F_L(a, z) = \sum_{i \geq 0} F_i(a; L) z^{i+1-\mu(L)}.$$

Of course, $F_i(a; L)$ ’s are in $Z[a^{\pm 1}]$. It is also known that $P_0(a^{-1}; L) = F_0(a; L)$ [20], while Skein and Kauffman polynomials are entirely independent.

The main result in this Chapter will be

THEOREM 3.1. *Suppose K is a knot with period r . Write*

$$F_0(a; K) + (a - a^{-1})F_1(a; K) = \sum_n \alpha_n a^n,$$

where α_n ’s are integers. Then, $\alpha_n \equiv 0 \pmod{r}$ unless $n \equiv 0 \pmod{r}$.

Furthermore, for $1 \leq i \leq \frac{r-3}{2}$, we have

$$\begin{aligned} F_{2i-1}(a; K) + (a - a^{-1})F_{2i}(a; K) &\equiv f_{2i}(a - a^{-1})F_0(a; K) \pmod{r}, \\ F_{2i}(a; K) + (a - a^{-1})F_{2i+1}(a; K) &\equiv 0 \pmod{r}, \end{aligned}$$

where f_{2i} 's, $2 \leq 2i \leq r-3$, are appropriate integers.

Taking the polynomial

$$(1 + \frac{a - a^{-1}}{z})F_K(a, z)$$

rather than $F_K(a, z)$, we will better understand what Theorem 3.1 implies.

Here, we shall present an example. Let K be a knot with period 5, made with 5 copies of

Then,

$$\begin{aligned} F_K(a, z) = & 7a^{-2} - 18 + 22a^2 - 15a^4 + 5a^6 \\ & + (2a^{-1} - 2a + 10a^5 - 15a^7 + 5a^9)z \\ & + (56a^{-2} - 202 + 301a^2 - 245a^4 + 95a^6 + 10a^8 - 15a^{10})z^2 \\ & + (76a^{-1} - 181a + 155a^3 - 5a^5 - 95a^7 + 45a^9 + 5a^{11})z^3 \\ & + \dots \end{aligned}$$

So, we have

$$\begin{aligned} (a - a^{-1})F_0(a; K) &\equiv 2a^3 - 2a^{-3} \pmod{5} \\ F_0(a; K) + (a - a^{-1})F_1(a; K) &\equiv 1 \pmod{5}, \\ F_1(a; K) + (a - a^{-1})F_2(a; K) &\equiv a^3 - a^{-3} \pmod{5} \\ F_2(a; K) + (a - a^{-1})F_3(a; K) &\equiv 0 \pmod{5}. \end{aligned}$$

Required regularities are apparent.

The proof of Theorem 3.1 will be given in section 3.6. The essential part of the proof of Theorem 3.1 will be, as in Skein polynomials, to compute Kauffman polynomials of torus knots and links. To do this in sections 3.4 and 3.5, we shall prepare some remarkable results from the view point of representation theory in sections 3.2 and 3.3.

3.2. BRAUER'S ALGEBRAS.

We will first define "Brauer's algebras" $D_n(x)$ over $\mathbb{C}$, where x is a complex but not an integer. For $n > 0$, a linear basis of $D_n(x)$ is given by graphs with n edges and $2n$ vertices, arranged in 2 lines of n vertices each. In these graphs, each edge belongs to exactly 2 vertices and each vertex belongs to exactly one edge. So the dimension of $D_n(x)$ is

$$1 \cdot 3 \cdot 5 \cdots (2n - 1).$$

Then, the product $\xi\eta$ of 2 graphs ξ and η can be defined similarly as braids. When $\xi\eta$ produces some closed components, define $\xi\eta = x^d\zeta$, where d denotes the number of such closed components and ζ denotes this graph without closed components.

It will be clear that Brauer's algebra $D_n(x)$ is generated by graphs $h_1, h_2, \dots, h_{n-1}$ and $e_1, e_2, \dots, e_{n-1}$ presented below.

e_i

h_i

We now describe the structure of $D_n(x)$. Let Γ_n denote the set of all Young diagrams with $k \leq n$ nodes such that $k \geq 0$ and $n - k$ is even.

THEOREM 3.2[42]. *Brauer's algebra $D_n(x)$ is semisimple. It decomposes into a direct sum of full matrix algebras $D_{n,\lambda}$, where $\lambda \in \Gamma_n$.*

To see this graphically, we shall prepare several notations. Let ν be the greatest integer which does not exceed $n/2$. For $0 \leq i \leq \nu$, we define $\bar{V}_{n,i}$ as a subalgebra of $D_n(x)$ generated by graphs with $k \geq 2i$ "horizontal" edges, where we call an edge *horizontal* if it joins 2 vertices in the same row. Then, it follows

$$\bar{V}_{n,0} \supset \bar{V}_{n,1} \supset \cdots \supset \bar{V}_{n,\nu}.$$

So we define

$$V_{n,i} = \bar{V}_{n,i} / \bar{V}_{n,i+1}$$

for $0 \leq i \leq \nu - 1$ and $V_{n,\nu} = \bar{V}_{n,\nu}$. Since $D_n(x)$ is semisimple, each canonical projection

$$\pi_{n,i} : \bar{V}_{n,i} \rightarrow V_{n,i}$$

splits, and we have

$$D_n(x) \cong V_{n,0} \oplus V_{n,1} \oplus \cdots \oplus V_{n,\nu}.$$

Note that each $V_{n,i}$ has a linear basis given by graphs with $2i$ horizontal edges. We denote the set of all graphs with $2i$ horizontal edges by $\mathcal{D}_{n,i}$. Unfortunately, $V_{n,i}$ are still nonsimple. In fact, from the construction of $D_{n,\lambda}$ in [], we can observe

$$V_{n,i} \cong \bigoplus_{\lambda \in \Lambda_{n-2i}} D_{n,\lambda},$$

where we denote by $\Lambda_{n-2i} \subset \Gamma_n$ the set of Young diagrams with $n - 2i$ nodes. Hence we now study the structure of each $V_{n,i}$.

For graphs $\xi, \eta \in \mathcal{D}_{n,i}$, we say ξ is *left equivalent* or *right equivalent* to η if horizontal edges joining vertices in the lower or upper row of ξ are arranged in the same manner as ones in η respectively and we say ξ is *equivalent* to η if ξ is left and right equivalent to η . For example, the following 2 graphs are left equivalent but not equivalent.

It is clear that the number of left equivalence classes in $\mathcal{D}_{n,i}$ equals

$$N_{n,i} = \begin{cases} 1 & \text{if } i = 0, \\ \binom{n}{2} \binom{n-2}{2} \cdots \binom{n-2i+2}{2} / i! & \text{if } i \neq 0. \end{cases}$$

and the number of equivalence classes of $\mathcal{D}_{n,i}$ equals $N_{n,i}^2$. Let $\mathcal{E}_{j,k}$'s, $0 \leq j, k \leq N_{n,i}$, denote the equivalence classes of $\mathcal{D}_{n,i}$ with all the elements in $\mathcal{E}_{j,1} \cup \mathcal{E}_{j,2} \cup \cdots \cup \mathcal{E}_{j,N_i}$ left equivalent. Then, the elements of each $\mathcal{E}_{j,k}$ are in one to one correspondence with ones of symmetric group S_{n-2i} . Hence we shall identify the vector space $E_{j,k}$ over $\mathbb{C}$ generated by $\mathcal{E}_{j,k}$ with $\mathbb{C}S_{n-2i}$. Let $E_{j,k,\lambda}$ and $F_{j,k,\lambda}$ be subspaces of $E_{j,k}$ which are identified with simple and irreducible components of $\mathbb{C}S_{n-2i}$ corresponding to $\lambda \in \Lambda_{n-2i}$ and we define

$$E_{n,\lambda} = \pi_{n,i} \left(\sum_{1 \leq j,k \leq N_{n,i}} E_{j,k,\lambda} \right),$$

$$F_{n,\lambda} = \pi_{n,i} \left(\sum_{1 \leq k \leq N_{n,i}} F_{1,k,\lambda} \right).$$

Then, a graphical observation will reveal that each $E_{n,\lambda}$ is a simple $D_n(x)$ module, that is,

$$V_{n,i} = \bigoplus_{\lambda \in \Lambda_{n-2i}} E_{n,\lambda}$$

and an irreducible component of $E_{n,\lambda}$ is given by $F_{n,\lambda}$. Counting dimensions, we can observe $D_{n,\lambda} \cong E_{n,\lambda}$. Let $\psi_{n,\lambda}$ denote the irreducible character of $D_n(x)$ corresponding to $E_{n,\lambda} \cong D_{n,\lambda}$. We will later need the following proposition.

PROPOSITION 3.3. *Suppose m and n are relatively prime. Then, we have*

$$\psi_{n,\lambda}(h_1 h_2 \dots h_{n-1})^m = \begin{cases} (-1)^\gamma & \text{if } \lambda \in \Lambda'_n, \\ 1 & \text{if } \lambda = \phi, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. An irreducible representation of $(h_1 h_2 \dots h_{n-1})^m$ corresponding to $\lambda \in \Gamma_n$ can be constructed from the action of $(h_1 h_2 \dots h_{n-1})^m$ on the vector space $F_{n,\lambda}$. Let $\mathcal{F}_{k,\lambda}$ denote a basis of $F_{1,k,\lambda}$ for $\lambda \in \Lambda_{n-2i}$. Then, a basis of $F_{n,\lambda}$ is given by

$$\mathcal{F}_{1,\lambda} \cup \mathcal{F}_{2,\lambda} \cup \dots \cup \mathcal{F}_{N_i,\lambda}$$

and $\mathcal{F}_{k,\lambda} \cap \mathcal{F}_{l,\lambda} = \emptyset$ if $k \neq l$. Furthermore, if $v \in \mathcal{F}_{k,\lambda}$, then v should be represented as a linear combination of graphs in $\mathcal{E}_{1,k}$. Therefore, from a graphical observation, it is not so difficult to check

$$(h_1 h_2 \dots h_{n-1})^m v \notin \mathcal{F}_{k,\lambda}$$

if $i \neq 0$ or $n - 2i \neq 0$. In this case, diagonal elements in the representation matrix are all zeroes, and so

$$\psi_{n,\lambda}(h_1 h_2 \dots h_{n-1})^m = 0.$$

Suppose $n - 2i = 0$, that is, $\lambda = \phi$. We note that each $\mathcal{F}_{k,\lambda}$ consists of an element, say v_k , and we can also observe there uniquely exists k' , $1 \leq k' \leq N_i$, such that

$$(h_1 h_2 \dots h_{n-1})^m v_{k'} = v_{k'}$$

and

$$(h_1 h_2 \dots h_{n-1})^m v_k \notin \mathcal{F}_{k,\lambda}$$

if $k \neq k'$. Hence we have

$$\psi_{n,\lambda}(h_1 h_2 \dots h_{n-1})^m = 1.$$

We now suppose $i = 0$, that is, $\lambda \in \Lambda_n$. Then, each $\psi_{n,\lambda}$ coincides with the irreducible character of CS_n corresponding to $\lambda \in \Lambda_n$. Thus, we have Proposition 3.3. $\square$

3.3. BIRMAN-WENZL-MURAKAMI ALGEBRAS.

We now introduce "Birman-Wenzl-Murakami algebras" $C_n(\alpha, q)$. Let α and q be non-zero complex numbers. We further suppose q is not a root of unity. Then, for an integer $n \geq 2$, $C_n(\alpha, q)$ will be defined as an algebra over $\mathbf{C}$ with generators

$$1, \tau_1^{\pm 1}, \tau_2^{\pm 1}, \dots, \tau_{n-1}^{\pm 1}, \epsilon_1, \epsilon_2, \dots, \epsilon_{n-1}$$

and relations

$$\begin{aligned} \tau_i \tau_{i+1} \tau_i &= \tau_{i+1} \tau_i \tau_{i+1}, & \epsilon_i \epsilon_{i \pm 1} \epsilon_i &= \epsilon_i, \\ \tau_i^{\pm 1} \epsilon_{i+1} \epsilon_i &= \tau_{i+1}^{\mp 1} \epsilon_i, & \epsilon_i \epsilon_{i+1} \tau_i^{\pm 1} &= \epsilon_i \tau_{i+1}^{\mp 1}, \\ \tau_i^{\pm 1} \epsilon_{i-1} \epsilon_i &= \tau_{i-1}^{\mp 1} \epsilon_i, & \epsilon_i \epsilon_{i-1} \tau_i^{\pm 1} &= \epsilon_i \tau_{i-1}^{\mp 1}, \\ \tau_i \tau_j &= \tau_j \tau_i, & \epsilon_i \tau_j &= \epsilon_j \tau_i, & \epsilon_i \epsilon_j &= \epsilon_j \epsilon_i & \text{ if } |i - j| > 1, \\ \tau_i \tau_i^{-1} &= \tau_i^{-1} \tau_i = 1, & \tau_i \epsilon_i &= \epsilon_i \tau_i = (-\alpha^2 q)^{-1} \epsilon_i, \\ \tau_i - \tau_i^{-1} &= (q - q^{-1})(1 - \epsilon_i). \end{aligned}$$

THEOREM 3.4[3,28]. *Birman-Wenzl-Murakami algebra $C_n(\alpha, q)$ is semisimple. It decomposes into a direct sum of full matrix algebras which are parametrized by Γ_n . By replacing $\sqrt{-q^{x-2}}$ for α and taking a limit $q \rightarrow 1$, $C_n(\alpha, q)$ coincides with Brauer's algebra $D_n(x)$.*

Let $\rho_{n,\lambda}$ denote an irreducible representation of $C_n(\alpha, q)$ corresponding to $\lambda \in \Gamma_n$, which is explicitly given in [28]. From the definition, the dimension $d_{n,\lambda}$ of $\rho_{n,\lambda}$ equals the number of paths of order n from λ to $\phi \in \Lambda_0$. Recall $u_{n,\lambda}$ and $v_{n,\lambda}$ denote the numbers of paths of order $n-2$ from λ to ϕ and to ϕ . Furthermore, we define $w_{n,\lambda}$ as the number of paths of order $n-2$ from λ to ϕ . Clearly,

$$d_{n,\lambda} = u_{n,\lambda} + v_{n,\lambda} + w_{n,\lambda}.$$

We begin with the following proposition.

PROPOSITION 3.5. *The determinant of $\rho_{n,\lambda}(\tau_i)$ is given by*

$$\Delta_{n,\lambda} = (-1)^{v_{n,\lambda}} q^{u_{n,\lambda} - v_{n,\lambda}} (-\alpha^2 q)^{-w_{n,\lambda}}.$$

Proof. All the τ_i 's are conjugate in $C_n(\alpha, q)$, which means that all the $\rho_{n,\lambda}(\tau_i)$'s have the same determinant. So it suffices to calculate the determinant of $\rho_{n,\lambda}(\tau_1)$. From the definition[28], the matrix $\rho_{n,\lambda}(\tau_1)$ is diagonal and its diagonal elements consist of q , $-q^{-1}$ and $-\alpha^{-2}q^{-1}$ with multiplicities $u_{n,\lambda}$, $v_{n,\lambda}$ and $w_{n,\lambda}$ respectively. This completes the proof of Proposition 3.5. $\square$

We can evaluate $u_{n,\lambda}$, $v_{n,\lambda}$ and $w_{n,\lambda}$ for certain λ .

PROPOSITION 3.6. For $\lambda \in \Lambda'_n$, we have

$$u_{n,\lambda} = \binom{\beta + \gamma - 1}{\beta - 1}, \quad v_{n,\lambda} = \binom{\beta + \gamma - 1}{\gamma - 1}, \quad w_{n,\lambda} = 0$$

and hence

$$d_{n,\lambda} = \binom{\beta + \gamma}{\gamma}.$$

For $\phi \in \Lambda_0$, we have

$$\begin{aligned} u_{n,\phi} &= v_{n,\phi} \\ w_{n,\phi} &= \binom{n-2}{2} \binom{n-4}{2} \cdots \binom{2}{2} / (\nu - 1)! \\ d_{n,\phi} &= \binom{n}{2} \binom{n-2}{2} \cdots \binom{2}{2} / \nu!, \end{aligned}$$

where $n = 2\nu$.

Proof. We shall prove the last 2 equalities. Theorem 3.4 implies that $d_{n,\phi}$ coincides with the dimension $N_{n,\nu}$ of $F_{n,\phi}$. Furthermore, from the definition, the equality

$$w_{n,\phi} = d_{n-2,\phi}$$

is obvious. Hence we can deduce Proposition 3.6. $\square$

Let $\chi_{n,\lambda}$'s denote characters of $\rho_{n,\lambda}$'s.

PROPOSITION 3.7.

$$\chi_{n,\lambda}(\tau_1 \tau_2 \cdots \tau_{n-1})^{mn} = d_{n,\lambda} \Delta_{n,\lambda}^{mn(n-1)/d_{n,\lambda}}.$$

In particular, for $\lambda \in \Lambda'_n$, we have

$$\chi_{n,\lambda}(\tau_1 \tau_2 \cdots \tau_{n-1})^{mn} = \binom{\beta + \gamma}{\gamma} q^{(\beta - \gamma)mn}.$$

Proof. As $(\tau_1 \tau_2 \cdots \tau_{n-1})^{mn}$ is in the center of $C_n(\alpha, q)$ and $\rho_{n,\lambda}$ is an irreducible representation, $\rho_{n,\lambda}(\tau_1 \tau_2 \cdots \tau_{n-1})^{mn}$ is scalar. Its determinant is given by

$$\Delta_{n,\lambda}^{mn(n-1)}$$

using Proposition 3.5 and so, we have

$$\rho_{n,\lambda}(\tau_1 \tau_2 \cdots \tau_{n-1})^{mn} = \Delta_{n,\lambda}^{mn(n-1)/d_{n,\lambda}}.$$

Proposition 3.7 now follows immediately. $\square$

PROPOSITION 3.8. Suppose m and n are relatively prime. Then we have

$$\chi_{n,\lambda}(\tau_1 \tau_2 \dots \tau_{n-1})^m = \begin{cases} (-1)^\gamma q^{(\beta-\gamma)m} & \text{if } \lambda \in \Lambda'_n \\ (-\alpha^2 q)^{-m} & \text{if } \lambda = \phi, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. As $\rho_{n,\lambda}(\tau_1 \tau_2 \dots \tau_{n-1})^{mn}$ is scalar, $\rho_{n,\lambda}(\tau_1 \tau_2 \dots \tau_{n-1})^m$ is diagonalizable. If we diagonalize it, the diagonal elements must be of the form

$$\omega_{\lambda,i}(q^{u_{n,\lambda}-v_{n,\lambda}}(-\alpha^2 q)^{-w_{n,\lambda}})^{m(n-1)/d_{n,\lambda}},$$

where

$$\omega_{\lambda,1}, \omega_{\lambda,2}, \dots, \omega_{\lambda,d_{n,\lambda}}$$

are roots of unity. By Theorem 3.4 and Proposition 3.5, we can observe

$$\psi_{n,\lambda}(h_1 h_2 \dots h_{n-1})^m = \omega_{\lambda,1} + \omega_{\lambda,2} + \dots + \omega_{\lambda,d_{n,\lambda}}.$$

Now, Proposition 3.8 follows from Propositions 3.3 and 3.6. $\square$

Propositions 3.7 and 3.8 have a connection with Kauffman polynomials of torus knots and links via the character formula of this polynomial established by J. Murakami, which we now describe.

For $\lambda \in \Gamma_n$, we also denote by $\rho_{n,\lambda}$ an irreducible representation of B_n which arises from $\rho_{n,\lambda}$ with an obvious homomorphism $\sigma_i \rightarrow \tau_i$. Characters of $\rho_{n,\lambda}$'s, for simplicity, will be also denoted by $\chi_{n,\lambda}$'s. As noted in Chapter 2, for $\xi \in B_n$, its closure $\hat{\xi}$ represents an oriented link L in S^3 . Suppose $\lambda \in \Gamma_n$ have λ_i nodes in its i -th row and $\tilde{\lambda}_j$ nodes in its j -th column. Then, we define

$$G_\lambda = \prod_{i=1}^{\tilde{\lambda}_1} g_\lambda(i) \prod_{j=1}^{\lambda_i} \frac{[j-i; 1]}{[\lambda_i + \tilde{\lambda}_j - i - j + 1]},$$

where

$$[k; m] = \alpha^m q^k - \alpha^{-m} q^{-k}$$

and

$$g_\lambda(i) = \begin{cases} \frac{\prod_{j=i}^{\lambda_i+i-1} [\lambda_i + \lambda_j - i - j + 2; 2]}{\prod_{j=1}^{\lambda_i} [j - i + 1; 1]} & \text{if } \lambda_i + i > \tilde{\lambda}_1, \\ \frac{\prod_{j=i}^{\tilde{\lambda}_1} [\lambda_i + \lambda_j - i - j + 2; 2]}{\prod_{j=1}^{\lambda_i} [j - i + 1; 1] \prod_{j=1}^{\tilde{\lambda}_1 - \lambda_i - i + 1} [3 - 2i - j; 2]} & \text{if } \lambda_i + i \leq \tilde{\lambda}_1. \end{cases}$$

THEOREM 3.9[28]. The Kauffman polynomial $F_L(a, z)$ of L with $a = -\alpha^2 q$, $z = [1]$ is given by

$$F_L(-\alpha^2 q, [1]) = (-1)^n (-\alpha^2 q)^{-\epsilon(\xi)} \frac{[1]}{[1] - [1; 2]} \sum_{\lambda \in \Gamma_n} G_\lambda \chi_{n, \lambda}(\xi).$$

It seems so complicated to calculate G_λ . However, we can do for certain λ .

PROPOSITION 3.10. For $\lambda \in \Lambda'_n$,

$$G_\lambda = \frac{\prod_{i=-\gamma}^{\beta} [i+1; 2]}{[\beta]! [\gamma]! [n]} - \frac{\prod_{i=-\gamma}^{\beta} [i+1; 2]}{[\beta]! [\gamma]! [\beta - \gamma + 1; 2]}.$$

Proof. Note that $\tilde{\lambda}_1 = \gamma + 1$. From the definition,

$$\begin{aligned} \prod_{i=1}^{\gamma+1} g_\lambda(i) &= \frac{[2\beta + 2; 2] \prod_{i=-\gamma}^{\beta} [i+1; 2]}{[\beta - \gamma + 1; 2] \prod_{i=-\gamma}^{\beta} [i+1; 1]}, \\ \prod_{i=1}^{\gamma+1} \prod_{j=1}^{\lambda_i} \frac{[j-i; 1]}{[\lambda_i + \tilde{\lambda}_j - i - j + 1]} &= \frac{\prod_{i=-\gamma}^{\beta} [i; 1]}{[\beta]! [\gamma]! [n]}. \end{aligned}$$

Thus, Proposition 3.10 follows immediately. We shall omit details. $\square$

Now, we can explicitly write down Kauffman polynomials of torus knots.

THEOREM 3.11. Let K be the (m, n) torus knot, where m and n are relatively prime and n is positive. The Kauffman polynomial of K is given by

$$a^{m(n-1)} \left(1 + \frac{a - a^{-1}}{[1]}\right) F_K(a, [1]) = \Sigma_a(m, n) + \bar{\Sigma}_a(m, n) + \begin{cases} 0 & \text{if } n \text{ is odd} \\ a^{-m} & \text{if } n \text{ is even,} \end{cases}$$

where

$$\Sigma_a(m, n) = \sum_{\beta + \gamma + 1 = n} (-1)^\gamma q^{(\beta - \gamma)m} \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]! [\gamma]! [n]}$$

and

$$\bar{\Sigma}_a(m, n) = \sum_{\beta + \gamma + 1 = n} (-1)^\gamma q^{(\beta - \gamma)m} \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]! [\gamma]! (aq^{\beta - \gamma} - a^{-1}q^{\gamma - \beta})}.$$

Proof. Since K is represented as the closure of

$$(\sigma_1 \sigma_2 \dots \sigma_{n-1})^m,$$

Theorem 3.11 follows from Proposition 3.8, Theorem 3.9 and Proposition 3.10 by making $a = -\alpha^2 q$. $\square$

This result is analogous to Jones' result concerning Skein polynomials of torus knots described in Theorem 2.12. We can easily observe

PROPOSITION 3.12.

$$\Sigma_a(m, n) = a^{m(n-1)} \frac{a - a^{-1}}{[1]} P_K(a^{-1}, [1]).$$

If $m = \pm 1$, then K is trivial and $P_K(v, z) = F_K(a, z) = 1$. Using Theorem 3.11 and Proposition 3.12, we can deduce the following 2 equalities which we will use later.

First, Proposition 3.12 tells us

COROLLARY 3.13.

$$\Sigma_a(\pm 1, n) = a^{\pm(n-1)} \frac{a - a^{-1}}{[1]}.$$

Combining this with Theorem 3.11,

COROLLARY 3.14.

$$\bar{\Sigma}_a(\pm 1, n) = \begin{cases} a^{\pm(n-1)} & \text{if } n \text{ is odd,} \\ a^{\pm(n-1)} - a^{\mp 1} & \text{if } n \text{ is even.} \end{cases}$$

We shall further prepare 2 identities.

PROPOSITION 3.15.

$$\Sigma_a(0, n) = \frac{a^n - a^{-n}}{[n]}.$$

Proof. Analogous to Proposition 2.13. $\square$

PROPOSITION 3.16.

$$\bar{\Sigma}_a(0, n) = \begin{cases} 1 & \text{if } n \text{ is odd,} \\ 0 & \text{if } n \text{ is even.} \end{cases}$$

Proof. It is easy to check that Proposition 3.16 holds for $n = 1, 2$. If $n > 2$,

$$\begin{aligned}
& (q^{n-1} + q^{-n+1})\bar{\Sigma}_a(0, n) - \bar{\Sigma}_a(1, n) - \bar{\Sigma}_a(-1, n) \\
&= \sum_{\beta+\gamma+1=n} (-1)^\gamma (q^{n-1} + q^{-n+1} - q^{\beta-\gamma} - q^{\gamma-\beta}) \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} \\
&= \sum_{\beta+\gamma+1=n} (-1)^\gamma [\beta][\gamma] \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} \\
&= - \sum_{\beta+\gamma+1=n, \beta, \gamma \geq 1} (-1)^{\gamma-1} (aq^{-\gamma} - a^{-1}q^{\gamma})(aq^{\beta} - a^{-1}q^{-\beta}) \\
&\quad \times \frac{\prod_{i=-\gamma+1}^{\beta-1} (aq^i - a^{-1}q^{-i})}{[\beta-1]![\gamma-1]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} \\
&= - \sum_{\beta'+\gamma'+1=n-2} (-1)^{\gamma'} (a^2q^{\beta'-\gamma'} + a^{-2}q^{\gamma'-\beta'} - q^{n-1} - q^{-n+1}) \\
&\quad \times \frac{\prod_{i=-\gamma'}^{\beta'} (aq^i - a^{-1}q^{-i})}{[\beta']![\gamma']!(aq^{\beta'-\gamma'} - a^{-1}q^{\gamma'-\beta'})} \\
&= (q^{n-1} + q^{-n+1})\bar{\Sigma}_a(0, n-2) - a^2\bar{\Sigma}_a(1, n-2) - a^{-2}\bar{\Sigma}_a(-1, n-2).
\end{aligned}$$

Corollary 3.14 implies

$$\bar{\Sigma}_a(\pm 1, n-2) - a^{\pm 2}\bar{\Sigma}_a(\pm 1, n-2) = 0$$

and so, it follows

$$\bar{\Sigma}_a(0, n) = \bar{\Sigma}_a(0, n-2).$$

Thus, inductive argument will be available. $\square$

3.4. TORUS LINKS.

In this section, we will detect Kauffman polynomials of torus links. Recall that the (r, nr) torus link L_n can be represented as the closure of

$$(\sigma_1\sigma_2\cdots\sigma_{r-1})^{nr} \in B_r$$

and every component of L_n is unknotted. So, we can obtain the following as in Proposition 2.9.

PROPOSITION 3.17.

$$z^{r-3}(F_{L_n}(a, z) - a^{-nr(r-1)}F_{L_0}(a, z)) = \sum_{i \geq 2} F_i(a; L_n)z^{i-2}.$$

We now investigate the left hand side of Proposition 3.17.

PROPOSITION 3.18. The polynomial

$$[1]^{r-3}(F_{L_n}(a, [1]) - a^{-nr(r-1)}F_{L_0}(a, [1])) \in Z[a^{\pm 1}, q^{\pm 1}]$$

decomposes into

$$[\bar{r}] \frac{R_1(a, q)}{Q_1(a, q)} + r \frac{R_2(a, q)}{Q_2(a, q)},$$

where R_1, R_2, Q_1, Q_2 are in $Z[a^{\pm 1}, q^{\pm 1}]$ and neither $Q_1(a, 1)$ nor $Q_2(a, 1)$ is divisible by r .

Proof. From Proposition 3.7 and Theorem 3.9, we have

$$\begin{aligned} & [1]^{r-3}(F_{L_n}(a, [1]) - a^{-nr(r-1)}F_{L_0}(a, [1])) \\ &= a^{-nr(r-1)} \frac{[1]^{r-2}}{[1] + a - a^{-1}} \sum_{\lambda \in \Gamma_r} G_{\lambda} d_{r, \lambda} (\Delta_{r, \lambda}^{nr(r-1)/d_{r, \lambda}} - 1). \end{aligned}$$

We divide the summation in the right hand side into 2 parts. Proposition 3.18 will be proved at once if the following 2 claims are established. $\square$

CLAIM 3.18.1. The first term

$$[\bar{r}] \frac{R_1(a, q)}{Q_1(a, q)}$$

in Proposition 3.18 should be

$$a^{-nr(r-1)} \frac{[1]^{r-2}}{[1] + a - a^{-1}} \sum_{\lambda \in \Lambda'_r} G_{\lambda} d_{r, \lambda} (\Delta_{r, \lambda}^{nr(r-1)/d_{r, \lambda}} - 1).$$

Proof of Claim 3.18.1. Let λ and $\lambda^* \in \Lambda'_r$ be as in Proposition 2.10. If $\gamma = \beta$, the contribution of λ in the summation is 0. Hence we suppose $\gamma > \beta$. Then, the contribution of λ and λ^* is

$$\begin{aligned} & G_{\lambda} d_{r, \lambda} (\Delta_{r, \lambda}^{nr(r-1)} - 1) + G_{\lambda^*} d_{r, \lambda^*} (\Delta_{r, \lambda^*}^{nr(r-1)} - 1) \\ &= \frac{(q^{(\beta-\gamma)nr} - 1) \prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i}) + (q^{(\gamma-\beta)nr} - 1) \prod_{i=-\beta}^{\gamma} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]![r]} \\ &+ \frac{(q^{(\beta-\gamma)nr} - 1) \prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} + \frac{(q^{(\gamma-\beta)nr} - 1) \prod_{i=-\beta}^{\gamma} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\gamma-\beta} - a^{-1}q^{\beta-\gamma})}, \end{aligned}$$

Note that $\beta - \gamma$ is even and so, $q^{\pm(\beta-\gamma)nr} - 1$ is divisible by $[r]$. Hence, to prove Claim 3.18.1, it is sufficient to show

$$\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i}) \equiv \prod_{i=-\beta}^{\gamma} (aq^i - a^{-1}q^{-i}) \pmod{[r]}.$$

This will be analogous to Proposition 2.10. $\square$

CLAIM 3.18.2. The second term

$$r \frac{R_2(a, q)}{Q_2(a, q)}$$

in Proposition 3.18 should be

$$a^{-nr(r-1)} \frac{[1]^{r-2}}{[1] + a - a^{-1}} \sum_{\lambda \notin \Lambda'_r} G_\lambda d_{r,\lambda} (\Delta_{r,\lambda}^{nr(r-1)/d_{r,\lambda}} - 1).$$

Proof of Claim 3.18.2. If $\lambda \notin \Lambda_r$, then $d_{r,\lambda}$ should be divisible by $N_{r,i}$ for appropriate $i \neq 0$, and if $i \neq 0$, $N_{r,i}$ is divisible by r . Hence $d_{r,\lambda}$ is also divisible by r . If $\lambda \in \Lambda_r$ and $\notin \Lambda'_r$, we can observe $d_{r,\lambda}$ is also divisible by r using the hook length formula. These observations enable us to establish Claim 3.18.2. $\square$

LEMMA 3.19. For $0 \leq i \leq r$,

$$F_i(a; L_n) \equiv a^{-nr(r-1)} F_i(a; L_0) \pmod{r}.$$

Since

$$F_{L_0}(a, z) = \left(1 + \frac{a - a^{-1}}{z}\right)^{r-1},$$

it follows

$$(a - a^{-1}) F_0(a; L_n) \equiv a^{-nr(r-1)} (a^r - a^{-r}) \pmod{r}$$

and, for $0 \leq i \leq r-2$,

$$F_i(a; L_n) + (a - a^{-1}) F_{i+1}(a; L_n) \equiv 0 \pmod{r}.$$

Furthermore,

$$F_{r-1}(a; L_n) + (a - a^{-1}) F_r(a; L_n) \equiv 1 \pmod{r}.$$

Proof. Propositions 3.17 and 3.18 implies

$$\sum_{i \geq 2} (F_i(a; L_n) - a^{-nr(r-1)} F_i(a; L_0)) [1]^{i-2} = [\bar{r}] \frac{R_1(a, q)}{Q_1(a, q)} + r \frac{R_2(a, q)}{Q_2(a, q)}.$$

As in Chapter 2, derivate both sides with respect to q for i times, where $0 \leq i \leq r-2$. By taking a limit $q \rightarrow 1$, the right hand side always vanishes mod r and the left hand side will turn into a linear combination of

$$F_0(a; L_n) - a^{-nr(r-1)} F_0(a; L_0),$$

$$F_1(a; L_n) - a^{-nr(r-1)} F_1(a; L_0),$$

.....

$$F_i(a; L_n) - a^{-nr(r-1)} F_i(a; L_0).$$

Thus, we can prove Lemma 3.19 inductively. $\square$

3.5. TORUS KNOTS.

In this section, we now investigate Kauffman polynomials of torus knots. As in Chapter 2, we denote $(r, nr \pm k)$ torus knots by T_n and T_n^* respectively. Let us begin with the case $n = 0$ as a special one.

PROPOSITION 3.20.

$$\begin{aligned} & q^{(k-1)r} [\bar{k}] a^{r(k-1)} ([1] + a - a^{-1}) F_{T_0}(a, [1]) \\ & \equiv a^k - a^{-k} + \begin{cases} [k] & \text{if } k \text{ is odd} \\ [k] a^{-r} & \text{if } k \text{ is even} \end{cases} \pmod{[\bar{r}]} \end{aligned}$$

Proof. From Propositions 3.15, 3.16 and Theorem 3.11, the difference between the left and right hand sides should be

$$[k](q^{(k-1)r} \Sigma_a(r, k) - \Sigma_a(0, k) + q^{(k-1)r} \bar{\Sigma}_a(r, k) - \bar{\Sigma}_a(0, k)).$$

Here, $q^{(k-1)r} \Sigma_a(r, k) - \Sigma_a(0, k)$ is expanded as

$$\sum_{\beta+\gamma+1=k} (-1)^\gamma (q^{2\beta r} - 1) \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]![k]}.$$

Furthermore, $q^{(k-1)r} \bar{\Sigma}_a(r, k) - \bar{\Sigma}_a(0, k)$ is also expanded as

$$\sum_{\beta+\gamma+1=k} (-1)^\gamma (q^{2\beta r} - 1) \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})}.$$

Note that each $q^{2\beta r} - 1$ is divisible by $[r]$. Since r is prime and $k < r$, denominators can not have a factor $[r]$ and so,

$$q^{(k-1)r} \Sigma_a(r, k) - \Sigma_a(0, k) + q^{(k-1)r} \bar{\Sigma}_a(r, k) - \bar{\Sigma}_a(0, k) \equiv 0 \pmod{[\bar{r}]} \quad \square.$$

The following implies Theorem 3.1 holds for T_0 .

PROPOSITION 3.21.

$$F_0(a; T_0) + (a - a^{-1}) F_1(a; T_0) \equiv \begin{cases} a^{-r(k-1)} & \text{if } k \text{ is odd} \\ a^{-rk} & \text{if } k \text{ is even} \end{cases} \pmod{r}.$$

Furthermore, for $1 \leq i \leq \frac{r-3}{2}$, we have

$$\begin{aligned} F_{2i-1}(a; T_0) + (a - a^{-1}) F_{2i}(a; T_0) & \equiv f_{2i}(a - a^{-1}) F_0(a; T_0) \pmod{r}, \\ F_{2i}(a; T_0) + (a - a^{-1}) F_{2i+1}(a; T_0) & \equiv 0 \pmod{r}, \end{aligned}$$

where f_{2i} 's are appropriate integers.

Proof. By Proposition 3.20, we have

$$\begin{aligned} 0 \equiv & q^{(k-1)r} [\bar{k}] a^{r(k-1)} (a - a^{-1}) F_0(a; T_0) - a^k + a^{-k} \\ & + q^{(k-1)r} [\bar{k}] a^{r(k-1)} (F_0(a; T_0) + (a - a^{-1}) F_1(a; T_0) - \begin{cases} a^{-r(k-1)} & \text{if } k \text{ is odd} \\ a^{-rk} q^r & \text{if } k \text{ is even} \end{cases}) [1] \\ & + q^{(k-1)r} [\bar{k}] a^{r(k-1)} \sum_{i \geq 2} (F_{i-1}(a; T_0) + (a - a^{-1}) F_i(a; T_0)) [1]^i \pmod{[\bar{r}]}. \end{aligned}$$

Derivate both sides with respect to q for i times, where $1 \leq i \leq r-2$. By taking a limit $q \rightarrow 1$, the right hand side will vanish mod r and the left hand side will be a linear combination of

$$\begin{aligned} & a^{r(k-1)} (a - a^{-1}) F_0(a; T_0), \\ & a^{r(k-1)} (F_0(a; T_0) + (a - a^{-1}) F_1(a; T_0) - \begin{cases} a^{-r(k-1)} & \text{if } k \text{ is odd} \\ a^{-rk} & \text{if } k \text{ is even} \end{cases}), \\ & a^{r(k-1)} (F_1(a; T_0) + (a - a^{-1}) F_2(a; T_0)), \\ & \dots\dots\dots \\ & a^{r(k-1)} (F_{i-1}(a; T_0) + (a - a^{-1}) F_i(a; T_0)). \end{aligned}$$

From the property of Kauffman polynomials[17], every term must be an odd power of a in $F_{2j-1}(a; T_0) + (a - a^{-1}) F_{2j}(a; T_0)$ and even in $F_{2j}(a; T_0) + (a - a^{-1}) F_{2j+1}(a; T_0)$. This implies both linear combinations of

$$\begin{aligned} & a^{r(k-1)} (a - a^{-1}) F_0(a; T_0), \\ & a^{r(k-1)} (F_1(a; T_0) + (a - a^{-1}) F_2(a; T_0)), \\ & \dots\dots\dots \\ & a^{r(k-1)} (F_{2j-1}(a; T_0) + (a - a^{-1}) F_{2j}(a; T_0)). \end{aligned}$$

and of

$$\begin{aligned} & a^{r(k-1)} (F_0(a; T_0) + (a - a^{-1}) F_1(a; T_0) - \begin{cases} a^{-r(k-1)} & \text{if } k \text{ is odd} \\ a^{-rk} & \text{if } k \text{ is even} \end{cases}), \\ & a^{r(k-1)} (F_2(a; T_0) + (a - a^{-1}) F_3(a; T_0)), \\ & \dots\dots\dots \\ & a^{r(k-1)} (F_{2j}(a; T_0) + (a - a^{-1}) F_{2j+1}(a; T_0)). \end{aligned}$$

should vanish mod r . Thus, we can prove Proposition 3.21 inductively. $\square$

Now, let us consider $n = -1$ as another special case.

PROPOSITION 3.22.

$$F_0(a; T_{-1}) + (a - a^{-1})F_1(a; T_{-1}) \equiv \begin{cases} a^{r(k-r)} & \text{if } k \text{ is odd} \\ a^{r(k-r+1)} & \text{if } k \text{ is even} \end{cases} \pmod{r}.$$

Furthermore, for $1 \leq i \leq \frac{r-3}{2}$, we have

$$\begin{aligned} F_{2i-1}(a; T_{-1}) + (a - a^{-1})F_{2i}(a; T_{-1}) &\equiv f_{2i}(a - a^{-1})F_0(a; T_{-1}) \pmod{r}, \\ F_{2i}(a; T_{-1}) + (a - a^{-1})F_{2i+1}(a; T_{-1}) &\equiv 0 \pmod{r}, \end{aligned}$$

where f_{2i} 's are the same integers as in Proposition 3.21.

Before the proof, we should note

PROPOSITION 3.23.

$$F_i(a; T_n^*) = \begin{cases} F_i(a^{-1}; T_{-n}) & \text{if } i \text{ is even,} \\ -F_i(a^{-1}; T_{-n}) & \text{if } i \text{ is odd.} \end{cases}$$

Proof. Since T_n^* is the mirror image of T_{-n} , we have

$$F_{T_n^*}(a, z) = F_{T_{-n}}(a^{-1}, -z)$$

[17]. Now, Proposition 3.23 follows immediately. $\square$

Proof of Proposition 3.22. By Proposition 3.23, it suffices to show for T_1^* . As in Proposition 3.21, we have

$$\begin{aligned} 0 &\equiv q^{(r-k-1)r} [r - k] a^{r(r-k-1)} (a - a^{-1}) F_0(a; T_1^*) - a^{r-k} + a^{k-r} \\ &\quad + q^{(r-k-1)r} [r - k] a^{r(r-k-1)} \\ &\quad \times (F_0(a; T_1^*) + (a - a^{-1})F_1(a; T_1^*) - \begin{cases} a^{-r(r-k-1)} & \text{if } k \text{ is even} \\ a^{-r(r-k)} q^r & \text{if } k \text{ is odd} \end{cases}) [1] \\ &\quad + q^{(r-k-1)r} [r - k] a^{r(r-k-1)} \sum_{i \geq 2} (F_{i-1}(a; T_1^*) + (a - a^{-1})F_i(a; T_1^*)) [1]^i \pmod{[r]}. \end{aligned}$$

Derivating and taking a limit $q \rightarrow 1$, we have a linear combination of

$$\begin{aligned} &a^{-r(r-k-1)} (a - a^{-1}) F_0(a; T_1^*), \\ &a^{-r(r-k-1)} (F_0(a; T_1^*) + (a - a^{-1})F_1(a; T_1^*) - \begin{cases} a^{-r(r-k-1)} & \text{if } k \text{ is odd} \\ a^{-r(r-k)} & \text{if } k \text{ is even} \end{cases}), \\ &a^{r(r-k-1)} (F_1(a; T_1^*) + (a - a^{-1})F_2(a; T_1^*)), \\ &\dots\dots\dots \\ &a^{r(r-k-1)} (F_{i-1}(a; T_1^*) + (a - a^{-1})F_i(a; T_1^*)) \end{aligned}$$

with the same coefficients as in Proposition 3.21 because

$$[r - k] \equiv -q^r[k] \pmod{[r]}.$$

Thus, we obtain

$$F_0(a; T_1^*) + (a - a^{-1})F_1(a; T_1^*) \equiv \begin{cases} a^{-r(r-k-1)} & \text{if } k \text{ is even,} \\ a^{-r(r-k)} & \text{if } k \text{ is odd} \end{cases} \pmod{r}$$

and

$$\begin{aligned} F_{2i-1}(a; T_1^*) + (a - a^{-1})F_{2i}(a; T_1^*) &\equiv f_{2i}(a - a^{-1})F_0(a; T_1^*) \pmod{r}, \\ F_{2i}(a; T_1^*) + (a - a^{-1})F_{2i+1}(a; T_1^*) &\equiv 0 \pmod{r} \end{aligned}$$

for $1 \leq i \leq \frac{r-3}{2}$, where f_{2i} 's are same integers as in Proposition 3.21. $\square$

Furthermore, we have

PROPOSITION 3.24.

$$a^{r(r-1)}F_{T_{n+1}}(a, [1]) - 2F_{T_n}(a, [1]) + a^{-r(r-1)}F_{T_{n-1}}(a, [1]) \equiv 0 \pmod{[\bar{r}]}.$$

This implies

$$a^{r(r-1)}F_i(a; T_{n+1}) - 2F_i(a; T_n) + a^{-r(r-1)}F_i(a; T_{n-1}) \equiv 0 \pmod{r}$$

for $0 \leq i \leq r-2$.

Proof. By Theorem 3.11,

$$F_{T_n}(a, [1]) = a^{-(nr+k)(r-1)} \frac{[1]}{[1] + a - a^{-1}} (\Sigma_a(nr+k, r) + \bar{\Sigma}_a(nr+k, r)).$$

Therefore,

$$\begin{aligned} &a^{r(r-1)}F_{T_{n+1}}(a, [1]) - 2F_{T_n}(a, [1]) + a^{-r(r-1)}F_{T_{n-1}}(a, [1]) \\ &= a^{-(nr+k)(r-1)} \frac{[1]}{[1] + a - a^{-1}} \sum_{\beta+\gamma+1=r} (-1)^\gamma (q^{(\beta-\gamma)r} - 2 + q^{(\gamma-\beta)r}) q^{nr+k} G_\lambda \\ &\equiv 0 \pmod{[\bar{r}]} \end{aligned}$$

because

$$q^{(\beta-\gamma)r} - 2 + q^{(\gamma-\beta)r} \equiv 0 \pmod{[r]^2}.$$

Thus, we have Proposition 3.24. $\square$

We shall close this section with the following 2 Lemmas. By Propositions 3.21, 3.22 and 3.24, we have

LEMMA 3.25. Theorem 3.1 holds for torus knots T_n , where f_{2i} 's are the same integers as in Proposition 3.21.

From Proposition 3.23, we can observe

$$\begin{aligned} F_{2i-1}(a; T_n^*) + (a - a^{-1})F_{2i}(a; T_n^*) &= -(F_{2i-1}(a^{-1}; T_n) + (a - a^{-1})F_{2i}(a^{-1}; T_n)), \\ F_{2i}(a; T_n^*) + (a - a^{-1})F_{2i+1}(a; T_n^*) &= F_{2i}(a^{-1}; T_n) + (a - a^{-1})F_{2i+1}(a^{-1}; T_n). \end{aligned}$$

So, it follows

LEMMA 3.26. Theorem 3.1 holds for torus knots T_n^* , where f_{2i} 's are the same integers as in Proposition 3.21.

3.6. PERIODIC REDUCTIONS.

We shall write

$$\Lambda_D(a, z) = \sum_{i \geq 1 - \mu(D)} \Lambda_i(a; D) z^i$$

for a diagram D of a link, where $\mu(D)$ denotes the number of components of the link which D represents.

LEMMA 3.27. Let L be a link with r components, having period r with no component invariant under rotation. Write

$$(a - a^{-1})F_0(a; L) = \sum_n \alpha_n a^n.$$

Then, $\alpha_n \equiv 0 \pmod{r}$ unless $n \equiv 0 \pmod{r}$.

Furthermore, for $0 \leq i \leq r - 2$, we have

$$F_i(a; L) + (a - a^{-1})F_{i+1}(a; L) \equiv 0 \pmod{r}.$$

Proof. L has a diagram D having period r in a plane. Define $D_{+\dots+}$, $D_{-\dots-}$, $D_{0\dots 0}$ and $D_{\infty\dots\infty}$ be D with all the crossings in a periodic orbit simaltenously replaced by

respectively. As in Chapter 2, we have another congruence[34]

$$\Lambda_{D_{+\dots+}}(a, z) - \Lambda_{D_{-\dots-}}(a, z) \equiv z^r (\Lambda_{D_{0\dots 0}}(a, z) - \Lambda_{D_{\infty\dots\infty}}(a, z)) \pmod{r}.$$

Recall

$$\text{min-deg}_z \wedge_D (a, z) = 1 - \mu(D).$$

From the assumption of D , we can easily observe

$$\mu(D_{0\dots 0}) = 2, \quad \mu(D_{\infty\dots\infty}) = 2$$

or

$$\mu(D_{0\dots 0}) = 2r, \quad \mu(D_{\infty\dots\infty}) = r.$$

In the upper case, from the congruence,

$$\wedge_i(a; D_{+\dots+}) - \wedge_i(a; D_{-\dots-}) \equiv 0 \pmod{r}$$

for $0 \leq i < 2(r-1)$, that is,

$$a^r F_i(a; D_{+\dots+}) - a^{-r} F_i(a; D_{-\dots-}) \equiv 0 \pmod{r}.$$

In the lower case,

$$\wedge_i(a; D_{+\dots+}) - \wedge_i(a; D_{-\dots-}) \equiv \wedge_i(a; D_{0\dots 0}) \pmod{r}$$

for $0 \leq i < r$, that is

$$a^r F_i(a; D_{+\dots+}) - a^{-r} F_i(a; D_{-\dots-}) \equiv F_i(a; D_{0\dots 0}) \pmod{r}.$$

Case 3.27.1. D is a closed nr braid.

We shall prove Lemma 3.27 by induction on n in this case. As was observed in Lemma 2.21, a sequence of periodic reductions will reduce D into a diagram of the (r, nr) torus link. Therefore, it too suffices to show that Lemma 3.27 holds for $D_{+\dots+}$ if and only if it does for $D_{-\dots-}$ by virtue of Lemma 3.19. If $\mu(D_{0\dots 0}) = 2$, this clearly holds. We suppose $\mu(D_{0\dots 0}) = 2r$. Then, our assertion is true if Lemma 3.27 holds for $D_{0\dots 0}$. So we check $F_i(a; D_{0\dots 0})$'s for $0 \leq i \leq r-1$. Note that $D_{0\dots 0}$ consists of 2 link diagrams, say $D'_{0\dots 0}, D''_{0\dots 0}$, satisfying the assumption of Lemma 3.27. Let

$$D_{0\dots 0, +\dots+}, \quad D_{0\dots 0, -\dots-}, \quad D_{0\dots 0, 0\dots 0}, \quad D_{0\dots 0, \infty\dots\infty}$$

denote D with all the crossings in a periodic orbit of a crossing between $D'_{0\dots 0}$ and $D''_{0\dots 0}$ replaced as above. Then, it follows

$$\mu(D_{0\dots 0, 0\dots 0}) = \mu(D_{0\dots 0, \infty\dots\infty}) = r$$

and so,

$$\wedge_i(a; D_{0\dots 0, +\dots+}) - \wedge_i(a; D_{0\dots 0, -\dots-}) \equiv 0 \pmod{r}$$

for $0 \leq i < 2r$. That is,

$$a^r F_i(a; D_{0\dots 0, +\dots +}) - a^{-r} F_i(a; D_{0\dots 0, -\dots -}) \equiv 0 \pmod{r}.$$

Repeating this, we can make $D_{0\dots 0}$ into a disjoint union $D'_{0\dots 0} \cup D''_{0\dots 0}$. As a result,

$$F_i(a; D_{0\dots 0}) \equiv a^{-2\nu r} F_i(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \pmod{r}$$

for $0 \leq i < 2r$, where

$$\nu r = \text{lk}(D'_{0\dots 0}, D''_{0\dots 0}).$$

Since

$$F_{D'_{0\dots 0} \cup D''_{0\dots 0}}(a, z) = \left(1 + \frac{a - a^{-1}}{z}\right) F_{D'_{0\dots 0}}(a, z) F_{D''_{0\dots 0}}(a, z)$$

[17], we have

$$F_0(a; D'_{0\dots 0} \cup D''_{0\dots 0}) = (a - a^{-1}) F_0(a; D'_{0\dots 0}) F_0(a; D''_{0\dots 0})$$

and for $i \geq 1$,

$$\begin{aligned} & F_i(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \\ &= \sum_{j=0}^{i-1} F_j(a; D'_{0\dots 0}) F_{i-j-1}(a; D''_{0\dots 0}) \\ &\quad + (a - a^{-1}) \sum_{j=0}^i F_j(a; D'_{0\dots 0}) F_{i-j}(a; D''_{0\dots 0}) \\ &= \sum_{j=0}^{i-1} F_j(a; D'_{0\dots 0}) (F_{i-j-1}(a; D''_{0\dots 0}) + (a - a^{-1}) F_{i-j}(a; D''_{0\dots 0})) \\ &\quad + (a - a^{-1}) F_i(a; D'_{0\dots 0}) F_0(a; D''_{0\dots 0}). \end{aligned}$$

Using the induction hypothesis, the first statement of Lemma 3.27 will be proved, and we further have

$$F_i(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \equiv (a - a^{-1}) F_i(a; D'_{0\dots 0}) F_0(a; D''_{0\dots 0}) \pmod{r}$$

for $0 \leq i \leq r - 1$. Consequently,

$$\begin{aligned} & F_i(a; D'_{0\dots 0} \cup D''_{0\dots 0}) + (a - a^{-1}) F_{i+1}(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \\ &\equiv (a - a^{-1}) F_0(a; D''_{0\dots 0}) (F_i(a; D'_{0\dots 0}) + (a - a^{-1}) F_{i+1}(a; D'_{0\dots 0})) \\ &\equiv 0 \pmod{r} \end{aligned}$$

will be deduced for $0 \leq i \leq r - 2$ also from the induction hypothesis. Thus, Lemma 3.27 holds for $D'_{0\dots 0} \cup D''_{0\dots 0}$ and so, for $D_{0\dots 0}$.

Case 3.27.2. Otherwise.

If D has no crossing point, then

$$F_D(a, z) = \left(1 + \frac{a - a^{-1}}{z}\right)^{r-1}$$

and so, Lemma 3.27 clearly holds. Now, as in the proof of Lemma 2.21, induction on the number of crossings in D will reduce this case into previous case, and will then complete the proof of Lemma 3.27. $\square$

Proof of Theorem 3.1. Suppose D is a diagram of a knot K with period r and k denotes the linking number between K and the axis of rotation. Define $D_{+...+}$, $D_{-...-}$, $D_{0...0}$ and $D_{\infty...\infty}$ as in Lemma 3.27. We can now observe

$$\mu(D_{0...0}) = 2, \quad \mu(D_{\infty...\infty}) = 1$$

or

$$\mu(D_{0...0}) = 2, \quad \mu(D_{\infty...\infty}) = r,$$

or

$$\mu(D_{0...0}) = r + 1, \quad \mu(D_{\infty...\infty}) = 1.$$

So, as in Lemma 3.27, the congruence tells us

$$\begin{aligned} & \Lambda_i(a; D_{+...+}) - \Lambda_i(a; D_{-...-}) \\ &= \begin{cases} 0 & \text{if } \mu(D_{0...0}) = 2 \text{ and } \mu(D_{\infty...\infty}) = 1 \\ \Lambda_{i-1}(a; D_{\infty...\infty}) & \text{if } \mu(D_{0...0}) = 2 \text{ and } \mu(D_{\infty...\infty}) = r \\ \Lambda_i(a; D_{0...0}) & \text{if } \mu(D_{0...0}) = r + 1. \end{cases} \end{aligned}$$

Case 3.1.1. D is a closed $nr \pm k$ braid, where we suppose $1 \leq k \leq r - 1$.

We shall prove Theorem 3.1 by induction on n . As in the proof of Theorem 2.2, a sequence of periodic reductions will reduce D into a diagram of the $(r, nr \pm k)$ torus knots and so, it suffices to show that Theorem 3.1 holds for $D_{+...+}$ if and only if it does for $D_{-...-}$ by virtue of Lemmas 3.25 and 3.26.

If $\mu(D_{0...0}) = 2$ and $\mu(D_{\infty...\infty}) = 1$, it is obvious. We suppose $\mu(D_{0...0}) = 2$ and $\mu(D_{\infty...\infty}) = r$. From the congruence, it follows

$$\begin{aligned} & (\Lambda_0(a; D_{+...+}) + (a - a^{-1}) \Lambda_1(a; D_{+...+})) \\ & - (\Lambda_0(a; D_{-...-}) + (a - a^{-1}) \Lambda_1(a; D_{-...-})) \\ & \equiv (a - a^{-1}) \Lambda_0(a; D_{\infty...\infty}) \pmod{r} \end{aligned}$$

and

$$\begin{aligned} & (\Lambda_i(a; D_{+...+}) + (a - a^{-1}) \Lambda_{i+1}(a; D_{+...+})) \\ & - (\Lambda_i(a; D_{-...-}) + (a - a^{-1}) \Lambda_{i+1}(a; D_{-...-})) \\ & \equiv \Lambda_{i-1}(D_{\infty...\infty}) + (a - a^{-1}) \Lambda_i(D_{\infty...\infty}) \pmod{r} \end{aligned}$$

for $1 \leq i \leq r-3$. Since $D_{\infty \dots \infty}$ should be a link with period r satisfying the assumption of Lemma 3.27, our assertion will follow from Lemma 3.27 and a fact

$$w(D_{+\dots+}) \equiv w(D_{-\dots-}) \equiv w(D_{\infty \dots \infty}) \equiv 0 \pmod{r}.$$

We now suppose $\mu(D_{0 \dots 0}) = r + 1$. From the congruence, our assertion will be proved if it is once established for $D_{0 \dots 0}$. Note that $D_{0 \dots 0}$ consists of a knot, say $D'_{0 \dots 0}$ and a link with r components, say $D''_{0 \dots 0}$. Define $D_{0 \dots 0, + \dots +}$, $D_{0 \dots 0, - \dots -}$, $D_{0 \dots 0, 0 \dots 0}$ and $D_{0 \dots 0, \infty \dots \infty}$ as in the proof of Lemma 3.27. Then, it will be easy to show

$$\mu(D_{0 \dots 0, 0 \dots 0}) = \mu(D_{0 \dots 0, \infty \dots \infty}) = 1.$$

Now, it follows

$$\Lambda_i(a; D_{0 \dots 0, + \dots +}) \equiv \Lambda_i(a; D_{0 \dots 0, - \dots -}) \pmod{r},$$

that is,

$$a^r F_i(a; D_{0 \dots 0, + \dots +}) \equiv a^{-r} F_i(a; D_{0 \dots 0, - \dots -}) \pmod{r}$$

for $0 \leq i < 2r$. Repeating periodic reductions, we can make $D_{0 \dots 0}$ into a disjoint union $D'_{0 \dots 0} \cup D''_{0 \dots 0}$ and so,

$$F_i(a; D_{0 \dots 0}) \equiv a^{-2\nu r} F_i(a; D'_{0 \dots 0} \cup D''_{0 \dots 0}) \pmod{r}$$

for $0 \leq i < 2r$. Since $D''_{0 \dots 0}$ satisfies the assumption of Lemma 3.27, as above, we have

$$F_i(a; D'_{0 \dots 0} \cup D''_{0 \dots 0}) \equiv (a - a^{-1}) F_0(a; D''_{0 \dots 0}) F_i(a; D'_{0 \dots 0}) \pmod{r}.$$

for $0 \leq i \leq r-1$. Note that induction hypothesis guarantees only for $i < r$, unfortunately.

Now our assertion has been reduced into $D_{0 \dots 0}$ and into $D'_{0 \dots 0}$. Since $D'_{0 \dots 0}$ has period r with mod r link number k against the axis of rotation, the induction hypothesis guarantees our assertion in this case and so Theorem 3.1 holds in Case 3.1.1.

Case 3.1.2. Otherwise.

As in Lemma 3.27, induction on the number of crossings in D will reduce this case into previous case. Now, we can complete the proof of Theorem 3.1. $\square$

4. A RELATIONSHIP BETWEEN SKEIN AND KAUFFMAN

4.1 MAIN THEOREM.

In this final Chapter, we shall discuss a curious correspondence between Skein and Kauffman polynomials for periodic knots. From the proof of Propositions 2.15 and 3.21, we will notice

$$p_{2i} = f_{2i}.$$

By the way, as was noted in previous Chapter,

$$P_0(a^{-1}; L) = F_0(a; L)$$

for any link L . Combining Theorems 2.2 and 3.1 with these facts, it follows

$$(a - a^{-1})P_i(a^{-1}; K) \equiv F_{i-1}(a; K) + (a - a^{-1})F_i(a; K) \pmod{r}$$

for

$$2 \leq i \leq r - 2$$

if K is a knot with period r . This is surprising because Skein and Kauffman polynomials are mutually independent, that is, one can not be a function of another. Of course, even for knots, such a correspondence will be almost broken in general case. However, more surprisingly, we can establish further relationship between them, that is,

$$(a - a^{-1})P_{r-1}(a^{-1}; K) \equiv F_{r-2}(a; K) + (a - a^{-1})F_{r-1}(a; K) \pmod{r}.$$

One may guess, perhaps, such a relationship will also appear in higher terms, but it seems the answer will be negative from an experiment.

We shall reformulate this curious relationship in terms of a new polynomial invariant derived from Skein and Kauffman polynomials. Define

$$J_L(a, z) = \left(1 + \frac{a - a^{-1}}{z}\right)F_L(a, z) - \frac{a - a^{-1}}{z}P_L(a^{-1}, z)$$

for a link L . Since $P_0(a^{-1}; L) = F_0(a; L)$, we can see

$$\min\text{-deg}_z J_L(a, z) = 1 - \mu(L).$$

So, as before, $J_L(a, z)$ can be decomposed as follows.

$$J_L(a, z) = \sum_{j \geq 0} J_j(a; L) z^{j+1-\mu(L)},$$

where $J_j(a; L)$'s are supposed to be in $Z[a^{\pm 1}]$. Note that

$$J_j(a; L) = F_j(a; L) + (a - a^{-1})F_{j+1}(a; L) - (a - a^{-1})P_{j+1}(a^{-1}; L).$$

We can now state the main result of this Chapter.

THEOREM 4.1. Suppose K is a knot with period r . Write

$$J_0(a; K) = \sum_n \alpha_n a^n,$$

where α_n 's are integers. Then, $\alpha_n \equiv 0 \pmod{r}$ unless $n \equiv 0 \pmod{r}$.

Furthermore, for $1 \leq j \leq r-2$, we have

$$J_j(a; K) \equiv 0 \pmod{r}.$$

The proof of Theorem 4.1 will be done in a direction analogous to Chapters 2 and 3. From an observation above, we only have to show

$$J_{r-2}(a; K) \equiv 0 \pmod{r}$$

and so, we shall only concern this $(r-2)$ -th term here. We will begin with torus knots as before.

4.2. TORUS KNOTS.

A formula of J for torus knots will be quite simple.

THEOREM 4.2. If K is the (m, n) torus knot,

$$a^{m(n-1)} J_K(a, [1]) = \bar{\Sigma}_a(m, n) + \begin{cases} 0 & \text{if } n \text{ is odd} \\ a^{-m} & \text{if } n \text{ is even.} \end{cases}$$

Proof of Theorem 4.2. From Theorem 2.12,

$$a^{m(n-1)} \frac{a - a^{-1}}{[1]} P_K(a^{-1}, [1]) = \Sigma_a(m, n)$$

and from Theorem 3.11,

$$a^{m(n-1)} \left(1 + \frac{a - a^{-1}}{[1]}\right) F_K(a, [1]) = \Sigma_a(m, n) + \bar{\Sigma}_a(m, n) + \begin{cases} 0 & \text{if } n \text{ is odd} \\ a^{-m} & \text{if } n \text{ is even.} \end{cases}$$

Now, Theorem 4.2 will be a corollary of 2 Theorems. $\square$

Let T_n and T_n^* be as before.

PROPOSITION 4.3.

$$J_{T_0}(a, z) \equiv \begin{cases} a^{(1-k)r} & \text{if } k \text{ is odd} \\ a^{-kr} q^r & \text{if } k \text{ is even} \end{cases} \pmod{[\bar{r}]}.$$

Proof of Proposition 4.3. By Theorem 4.2,

$$a^{(k-1)r} J_{T_0}(a, [1]) = \bar{\Sigma}_a(r, k) + \begin{cases} 0 & \text{if } k \text{ is odd} \\ a^{-r} & \text{if } k \text{ is even.} \end{cases}$$

On the other hand, from Corollary 3.16,

$$\bar{\Sigma}_a(0, k) = \begin{cases} 1 & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even} \end{cases}$$

and so,

$$\begin{aligned} & (aq)^{(k-1)r} J_{T_0}(a, [1]) - \begin{cases} 1 & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even} \end{cases} \\ &= q^{(k-1)r} \bar{\Sigma}_a(r, k) - \bar{\Sigma}_a(0, k) + \begin{cases} 0 & \text{if } k \text{ is odd} \\ a^{-r} & \text{if } k \text{ is even.} \end{cases} \\ &\equiv \sum_{\beta+\gamma+1=k} (-1)^\gamma (q^{2\beta r} - 1) \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} + \begin{cases} 0 & \text{if } k \text{ is odd} \\ a^{-r} & \text{if } k \text{ is even.} \end{cases} \\ &\equiv \begin{cases} 0 & \text{if } k \text{ is odd} \\ a^{-r} & \text{if } k \text{ is even.} \end{cases} \pmod{[\bar{r}]} \end{aligned}$$

Consequently,

$$J_{T_0}(a, z) \equiv \begin{cases} a^{(1-k)r} & \text{if } k \text{ is odd} \\ a^{-kr} q^r & \text{if } k \text{ is even} \end{cases} \pmod{[\bar{r}]} \quad \square$$

Using Proposition 4.3, we can prove

COROLLARY 4.4.

$$J_{r-2}(a; T_0) \equiv 0 \pmod{r}.$$

PROPOSITION 4.5.

$$a^{r(r-1)} J_{r-2}(a; T_{n+1}) \equiv J_{r-2}(a; T_n) \pmod{[\bar{r}]}.$$

Proof of Proposition 4.5. It follows from

$$\begin{aligned} & a^{r(r-1)} J_{T_{n+1}}(a, [1]) - J_{T_n}(a, [1]) \\ &= a^{-(nr+k)(r-1)} \sum_{\beta+\gamma+1=r} (-1)^\gamma (q^{(\beta-\gamma)r} - 1) q^{(\beta-\gamma)(nr+k)} \frac{\prod_{i=-\gamma}^{\beta} (aq^i - a^{-1}q^{-i})}{[\beta]![\gamma]!(aq^{\beta-\gamma} - a^{-1}q^{\gamma-\beta})} \\ &\equiv 0 \pmod{[\bar{r}]} \quad \square \end{aligned}$$

Consequently, we can obtain

LEMMA 4.6.

$$J_{r-2}(a; T_n) \equiv 0 \pmod{r}.$$

Similarly, we have

LEMMA 4.7.

$$J_{r-2}(a; T_n^*) \equiv 0 \pmod{r}.$$

4.3. PERIODIC REDUCTIONS.

From Lemmas 2.21 and 3.27, we already have

LEMMA 4.8. *Let L be a link with r components, having period r with no component invariant under rotation. Then, for $j \leq r - 2$,*

$$J_j(a; L) \equiv 0 \pmod{r}.$$

We can now prove Theorem 4.1.

Proof of Theorem 4.1. Suppose D is a diagram of a knot K with period r , and k denotes mod r linking number between K and the axis of rotation. Define $D_{+...+}$, $D_{-...-}$, $D_{0...0}$ and $D_{\infty...\infty}$ as before. We can again observe

$$\mu(D_{0...0}) = 2, \quad \mu(D_{\infty...\infty}) = 1$$

or

$$\mu(D_{0...0}) = 2, \quad \mu(D_{\infty...\infty}) = r,$$

or

$$\mu(D_{0...0}) = r + 1, \quad \mu(D_{\infty...\infty}) = 1.$$

Case 4.1.1. D is a closed $nr \pm k$ braid, where we suppose $1 \leq k \leq r - 1$.

We shall prove Theorem 4.1 by induction on n . Since a sequence of periodic reductions will reduce D into a diagram of the $(r, nr \pm k)$ torus knot, it suffices to show that Theorem 4.1 holds for $D_{+...+}$ if and only if it does for $D_{-...-}$ by virtue of Lemmas 4.6 and 4.7. If $\mu(D_{0...0}) = 2$ and $\mu(D_{\infty...\infty}) = 1$, we have

$$a^r P_{r-1}(a^{-1}; D_{+...+}) - a^{-r} P_{r-1}(a^{-1}; D_{-...-}) \equiv P_0(a^{-1}; D_{0...0}) \pmod{r}$$

and

$$a^r F_{r-2}(a; D_{+...+}) - a^{-r} F_{r-2}(a; D_{-...-}) \equiv 0 \pmod{r}$$

$$a^r F_{r-1}(a; D_{+...+}) - a^{-r} F_{r-1}(a; D_{-...-}) \equiv F_0(a; D_{0...0}) \pmod{r}.$$

So, it follows

$$\begin{aligned} a^r J_{r-2}(a; D_{+\dots+}) - a^{-r} J_{r-2}(a; D_{-\dots-}) &\equiv (a - a^{-1})(F_0(a; D_{0\dots 0}) - P_0(a^{-1}; D_{0\dots 0})) \\ &\equiv 0 \pmod{r}. \end{aligned}$$

We suppose $\mu(D_{0\dots 0}) = 2$ and $\mu(D_{\infty\dots\infty}) = r$. From the congruence, it follows

$$a^r P_{r-1}(a^{-1}; D_{+\dots+}) - a^{-r} P_{r-1}(a^{-1}; D_{-\dots-}) \equiv P_0(a^{-1}; D_{0\dots 0}) \pmod{r}$$

and

$$\begin{aligned} a^r F_{r-2}(a; D_{+\dots+}) - a^{-r} F_{r-2}(a; D_{-\dots-}) &\equiv a^{\kappa r} F_{r-3}(a; D_{\infty\dots\infty}) \pmod{r} \\ a^r F_{r-1}(a; D_{+\dots+}) - a^{-r} F_{r-1}(a; D_{-\dots-}) &\equiv F_0(a; D_{0\dots 0}) - a^{\kappa r} F_{r-2}(a; D_{\infty\dots\infty}) \pmod{r}, \end{aligned}$$

where κ will be some integer. Since $D_{\infty\dots\infty}$ should be a link with period r satisfying the assumption of Lemma 3.27, we have

$$\begin{aligned} &a^r (F_{r-2}(a; D_{+\dots+}) + (a - a^{-1})F_{r-1}(a; D_{+\dots+})) \\ &\quad - a^{-r} (F_{r-2}(a; D_{-\dots-}) + (a - a^{-1})F_{r-1}(a; D_{-\dots-})) \\ &\equiv (a - a^{-1})F_0(a; D_{0\dots 0}) + a^{\kappa r} (F_{r-3}(a; D_{\infty\dots\infty}) + (a - a^{-1})F_{r-2}(a; D_{\infty\dots\infty})) \\ &\equiv (a - a^{-1})F_0(a; D_{0\dots 0}) \pmod{r} \end{aligned}$$

and also,

$$a^r J_{r-2}(a; D_{+\dots+}) - a^{-r} J_{r-2}(a; D_{-\dots-}) \equiv 0 \pmod{r}.$$

We now suppose $\mu(D_{0\dots 0}) = r + 1$. Then,

$$a^r J_{r-2}(a; D_{+\dots+}) - a^{-r} J_{r-2}(a; D_{-\dots-}) \equiv J_{r-2}(a; D_{0\dots 0}) \pmod{r}.$$

From this, our assertion will be proved if once

$$J_{r-2}(a; D_{0\dots 0}) \equiv 0 \pmod{r}$$

is established. As was observed, $D_{0\dots 0}$ consists of a knot, say $D'_{0\dots 0}$ and a link with r components, say $D''_{0\dots 0}$. Define $D_{0\dots 0,+\dots+}$, $D_{0\dots 0,-\dots-}$, $D_{0\dots 0,0\dots 0}$ and $D_{0\dots 0,\infty\dots\infty}$ as in the proof of Lemma 3.27. Then, it will be easy to show

$$\mu(D_{0\dots 0,0\dots 0}) = \mu(D_{0\dots 0,\infty\dots\infty}) = 1.$$

Now, it follows

$$a^r J_{r-2}(a; D_{0\dots 0,+\dots+}) \equiv a^{-r} J_{r-2}(a; D_{0\dots 0,-\dots-}) \pmod{r}.$$

Repeating periodic reductions, we can make $D_{0\dots 0}$ into a disjoint union $D'_{0\dots 0} \cup D''_{0\dots 0}$ and so,

$$J_{r-2}(a; D_{0\dots 0}) \equiv a^{-2\nu r} J_{r-2}(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \pmod{r}.$$

By definition,

$$\begin{aligned}
& J_{D'_{0\dots 0} \cup D''_{0\dots 0}}(a, z) \\
&= (1 + \frac{a - a^{-1}}{z}) F_{D'_{0\dots 0} \cup D''_{0\dots 0}}(a, z) - \frac{a - a^{-1}}{z} P_{D'_{0\dots 0} \cup D''_{0\dots 0}}(a^{-1}, z) \\
&= (1 + \frac{a - a^{-1}}{z})^2 F_{D'_{0\dots 0}}(a, z) F_{D''_{0\dots 0}}(a, z) - (\frac{a - a^{-1}}{z})^2 P_{D'_{0\dots 0}}(a^{-1}, z) P_{D''_{0\dots 0}}(a^{-1}, z) \\
&= (1 + \frac{a - a^{-1}}{z}) F_{D''_{0\dots 0}}(a, z) J_{D'_{0\dots 0}}(a, z) + \frac{a - a^{-1}}{z} J_{D''_{0\dots 0}}(a, z) P_{D'_{0\dots 0}}(a^{-1}, z).
\end{aligned}$$

Since

$$J_{D'_{0\dots 0} \cup D''_{0\dots 0}}(a, z) = \sum_{j \geq 0} J_j(a; D'_{0\dots 0} \cup D''_{0\dots 0}) z^{j-r},$$

we can evaluate $J_{r-2}(a; D'_{0\dots 0} \cup D''_{0\dots 0})$ by looking at coefficients of z^{-2} . Since $D''_{0\dots 0}$ satisfies the assumption in Lemma 3.27,

$$\begin{aligned}
& (1 + \frac{a - a^{-1}}{z}) F_{D''_{0\dots 0}}(a, z) J_{D'_{0\dots 0}}(a, z) \\
&= \sum_{i \geq 0} (F_{i-1}(a; D''_{0\dots 0}) + (a - a^{-1}) F_i(a; D''_{0\dots 0})) z^{i-r} \times \sum_{j \geq 0} J_j(a; D'_{0\dots 0}) z^j \\
&= \sum_{i, j \geq 0} (F_{i-1}(a; D''_{0\dots 0}) + (a - a^{-1}) F_i(a; D''_{0\dots 0})) J_j(a; D'_{0\dots 0}) z^{i+j-r} \\
&\equiv (a - a^{-1}) F_0(a; D''_{0\dots 0}) \sum_{j \geq 0} J_j(a; D'_{0\dots 0}) z^{j-r} \\
&\quad + \sum_{i, j \geq 0} (F_{i+r-1}(a; D''_{0\dots 0}) + (a - a^{-1}) F_{i+r}(a; D''_{0\dots 0})) J_j(a; D'_{0\dots 0}) z^{i+j}.
\end{aligned}$$

Thus, z^{-2} has a coefficient

$$(a - a^{-1}) F_0(a; D''_{0\dots 0}) J_{r-2}(a; D'_{0\dots 0}).$$

Similarly, by Lemma 4.8,

$$\begin{aligned}
& \frac{a - a^{-1}}{z} J_{D''_{0\dots 0}}(a, z) P_{D'_{0\dots 0}}(a^{-1}, z) \\
&\equiv (a - a^{-1}) J_0(a; D''_{0\dots 0}) \sum_{i \geq 0} P_i(a; D'_{0\dots 0}) z^{i-r} \\
&\quad + (a - a^{-1}) \sum_{i, j \geq 0} J_{j+r-1}(a; D''_{0\dots 0}) P_i(a^{-1}; D'_{0\dots 0}) z^{i+j-1}
\end{aligned}$$

and so, z^{-2} has a coefficient

$$(a - a^{-1}) J_0(a; D'_{0\dots 0}) P_{r-2}(a; D''_{0\dots 0}) = 0.$$

Therefore, we obtain

$$J_{r-2}(a; D'_{0\dots 0} \cup D''_{0\dots 0}) \equiv (a - a^{-1})F_0(a; D''_{0\dots 0})J_{r-2}(a; D'_{0\dots 0}) \pmod{r}.$$

Now, our assertion has been reduced from $D_{0\dots 0}$ into $D'_{0\dots 0}$. Since $D'_{0\dots 0}$ has period r with mod r linking number k against the axis of rotation, the induction hypothesis guarantees our assertion in this case and so Theorem 4.1 holds in Case 4.1.1.

Case 4.1.2. Otherwise.

Induction on the number of crossings will reduce this case into previous case. Now, we can complete the proof of Theorem 4.1. $\square$

Finally, we shall close this paper with an example. Let K denote the $(3, 7)$ torus knot. Then, we have

$$\begin{aligned} \frac{a - a^{-1}}{z} P_K(a^{-1}, z) &= (5a^{17} - 21a^{15} + 28a^{13} - 12a^{11})z^{-1} \\ &\quad + (10a^{17} - 70a^{15} + 126a^{13} - 66a^{11})z \\ &\quad + (6a^{17} - 84a^{15} + 210a^{13} - 132a^{11})z^3 \\ &\quad + (a^{17} - 45a^{15} + 165a^{13} - 121a^{11})z^5 \\ &\quad + \dots, \\ (1 + \frac{a - a^{-1}}{z}) F_K(a, z) &= (5a^{17} - 21a^{15} + 28a^{13} - 12a^{11})z^{-1} \\ &\quad + (21a^{16} - 48a^{14} + 28a^{12}) \\ &\quad + (10a^{17} - 70a^{15} + 126a^{13} - 66a^{11})z \\ &\quad + (70a^{16} - 196a^{14} + 126a^{12})z^2 \\ &\quad + (6a^{17} - 84a^{15} + 210a^{13} - 132a^{11})z^3 \\ &\quad + (84a^{16} - 294a^{14} + 210a^{12})z^4 \\ &\quad + (a^{17} - 45a^{15} + 165a^{13} - 121a^{11})z^5 \\ &\quad + (45a^{16} - 210a^{14} + 165a^{12})z^6 \\ &\quad + \dots \end{aligned}$$

Thus, regularities described in Theorems 2.2 and 3.1 of course appear. Furthermore,

$$\begin{aligned} J_K(a, z) &= 21a^{16} - 48a^{14} + 28a^{12} \\ &\quad + (70a^{16} - 196a^{14} + 126a^{12})z^2 \\ &\quad + (84a^{16} - 294a^{14} + 210a^{12})z^4 \\ &\quad + (45a^{16} - 210a^{14} + 165a^{12})z^6 \\ &\quad + \dots \end{aligned}$$

Consequently, we have

$$J_0(a; K) \equiv a^{14} \pmod{7}$$

and

$$J_1(a; K) \equiv J_2(a; K) \equiv J_3(a; K) \equiv J_4(a; K) \equiv J_5(a; K) \equiv 0 \pmod{7}.$$

However, we shall note

$$J_6(a; K) \equiv 3a^{16} - 3a^{12} \pmod{7}.$$

Examples often give us valuable suggestions. Above all, torus knots has been nice, and essential ones for periodic knots, and I still guess there may be hidden features of polynomials of them. How can we discover such ones, and further criteria for periodic knots?

I sincerely hope results established here would give a clue not only for such works, but also for geometric characterization of new polynomial invariants.

REFERENCES

1. C. Adams, M. Hildebrand and J. Weeks, *Hyperbolic invariants of knots and links*, Trans. Amer. Math. Soc. **326** (1991), 1–56.
2. J.S. Birman, “Braids, links, and mapping class groups”, Ann. Math. Stud. 82, Princeton, 1976.
3. J.S. Birman and H. Wenzl, *Braids, link polynomials and a new algebra*, Trans. Amer. Math. Soc. **313** (1989), 249–273.
4. R. Brauer, *On algebras which are connected with the semisimple continuous groups*, Ann. Math. **38** (1937), 854–872.
5. G. Burde and H. Zieschang, “Knots”, de Gruyter Studies in Math. 5, Berlin, 1985.
6. J.F. Davis and C. Livingston, *Alexander polynomials of periodic knots*, Topology **30** (1991), 551–564.
7. ———, *Periodic knots, Smith theory, and Murasugi’s congruence*, L’Enseignement Math. (to appear).
8. P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K. Millett and A. Ocneanu, *A new polynomial invariant of knots and links*, Bull. Amer. Math. Soc. **12** (1985), 239–246.
9. C.McA. Gordon, R. Litherland and K. Murasugi, *Signatures of covering links*, Canad. J. Math. **33** (1981), 381–394.
10. J.A. Hillman, *Alexander ideals of links*, Lecture Note in Math. 895, Springer-Verlag, 1981.
11. P. Harpe, M. Kervaire and C. Weber, *On the Jones polynomial*, L’Enseignement Math. **32** (1986), 271–335.
12. V.F.R. Jones, *A polynomial invariant for knots via von Neumann algebra*, Bull. Amer. Math. Soc. **12** (1985), 103–111.
13. ———, *Hecke algebra representations of braid groups and link polynomials*, Ann. Math. **126** (1987), 335–388.
14. L.H. Kauffman, “Formal knot theory”, Math. Notes 30, Princeton, 1983.
15. ———, “On knots”, Ann. Math. Stud. 115, Princeton, 1987.
16. ———, *State models and the Jones polynomial*, Topology **26** (1987), 395–407.
17. ———, *An invariant of regular isotopy*, Trans. Amer. Math. Soc. **318** (1990), 417–471.
18. K. Kodama and M. Sakuma, *Symmetry groups of prime knots up to 10 crossings*, preprint.
19. W.B.R. Lickorish, *A relationship between link polynomials*, Math. Proc. Camb. Phil. Soc. **100** (1987), 109–112.
20. ———, *Polynomials for links*, Bull. London Math. Soc. **30** (1988), 558–588.
21. W.B.R. Lickorish and K.C. Millett, *A polynomial invariant of oriented links*, Topology **26** (1987), 107–141.
22. ———, *Some evaluations of link polynomials*, Comment. Math. Helvetici **61** (1986), 349–359.
23. U. Lüdicke, *Zyklische Knoten*, Archiv. Math. **32** (1979), 588–599.
24. ———, *9_{25} has no period 3*, C. R. Math. Rep. Acad. Sci. Canada **VI** (1984), 157–158.
25. J. Morgan and H. Bass, “The Smith conjecture”, Acad. Press, 1984.
26. J. Murakami, *The Kauffman polynomial of links and representation theory*, Osaka J. Math. **24**

- (1987), 745-758.
27. ———, *The parallel version of polynomial invariants of links*, Osaka J. Math. **26** (1989), 1-55.
  28. ———, *The representations of the q -analogue of Brauer's centralizer algebras and the Kauffman polynomial of links*, Publ. RIMS. Kyoto Univ. **26** (1990), 935-945.
  29. K. Murasugi, *On periodic knots*, Comment. Math. Helvetici **46** (1971), 162-174.
  30. ———, *On symmetries of knots*, Tsukuba J. Math. **4** (1980), 331-347.
  31. ———, *Jones polynomials and classical conjectures in knot theory*, Topology **26** (1987), 187-194.
  32. ———, *Jones polynomials and classical conjectures in knot theory II*, Math. Proc. Camb. Phil. Soc. **102** (1987), 317-318.
  33. ———, *Jones polynomials of periodic links*, Pacific J. Math. **131** (1988), 319-329.
  34. J.H. Przytycki, *On Murasugi's and Traczyk's criteria for periodic links*, Math. Ann. **283** (1989), 465-478.
  35. D. Rolfsen, "Knots and links", Publish or Perish Press, Berkeley, 1976.
  36. M. Sakuma, *On the polynomials of periodic links*, Math. Ann. **257** (1981), 487-494.
  37. H.N.V. Temperley and E. Lieb, *Relations between the "percolation" and "colouring" problem and other graph-theoretical problems associated with planar lattices*, Proc. Roy. Soc. London **322** (1971), 251-280.
  38. H.F. Trotter, *Periodic automorphisms of groups and knots*, Duke Math. J. **28** (1961), 553-557.
  39. P. Traczyk, *10_{101} has no period 7: a criterion for periodicity of links*, Proc. Amer. Math. Soc. **180** (1990), 845-846.
  40. ———, *A criterion for knots of period 3*, Topology and its appl. **36** (1990), 275-281.
  41. ———, *Periodic knots and the skein polynomial*, Invent. Math. **106** (1991), 73-84.
  42. H. Wenzl, *On the structure of Brauer's centralizer algebras*, Ann. Math. **128** (1988), 173-193.
  43. ———, *Hecke algebras of type A_n and subfactors*, Invent. Math. **92** (1988), 349-383.
  44. Y. Yokota, *The Jones polynomial of periodic knots*, Proc. Amer. Math. Soc. **113** (1991), 889-894.
  45. ———, *The skein polynomial of periodic knots*, Math. Ann. **291** (1991), 281-291.
  46. ———, *The Kauffman polynomial of periodic knots*, Topology (to appear).

A LIST OF PAPERS BY YOSHIYUKI YOKOTA

- [1] The Jones polynomial of periodic knots, Proc. Amer. Math. Soc. **113**, 889–894, 1991.
- [2] Twisting formulae of the Jones polynomial, Math. Proc. Camb. Phil. Soc. **110**, 473–482, 1991.
- [3] The skein polynomial of periodic knots, Math. Ann. **291**, 281–291, 1991.
- [4] Polynomial invariants of positive links, To appear in Topology.
- [5] The Kauffman polynomial of periodic knots, To appear in Topology.